

Lattice-valued functions

Eszter K. Horváth, Szeged

Co-authors: Branimir Šešelja, Andreja Tepavčević

Novi Sad, 2017, June 15. .

Let S be a nonempty set and L a complete lattice. Every mapping $\mu : S \rightarrow L$ is called a **lattice-valued** (L -valued) **function** on S .

Cut set (p -cut)

Let $p \in L$. A **cut set** of an L -valued function $\mu : S \rightarrow L$ (a p -cut) is a subset $\mu_p \subseteq S$ defined by:

$$x \in \mu_p \text{ if and only if } \mu(x) \geq p. \quad (1)$$

In other words, a p -cut of $\mu : S \rightarrow L$ is the inverse image of the principal filter $\uparrow p$, generated by $p \in L$:

$$\mu_p = \mu^{-1}(\uparrow p). \quad (2)$$

It is obvious that for every $p, q \in L$, $p \leq q$ implies $\mu_q \subseteq \mu_p$.

Cut set (p -cut)

Let $p \in L$. A **cut set** of an L -valued function $\mu : S \rightarrow L$ (a p -cut) is a subset $\mu_p \subseteq S$ defined by:

$$x \in \mu_p \text{ if and only if } \mu(x) \geq p. \quad (1)$$

In other words, a p -cut of $\mu : S \rightarrow L$ is the inverse image of the principal filter $\uparrow p$, generated by $p \in L$:

$$\mu_p = \mu^{-1}(\uparrow p). \quad (2)$$

It is obvious that for every $p, q \in L$, $p \leq q$ implies $\mu_q \subseteq \mu_p$.

Cut set (p -cut)

Let $p \in L$. A **cut set** of an L -valued function $\mu : S \rightarrow L$ (a p -cut) is a subset $\mu_p \subseteq S$ defined by:

$$x \in \mu_p \text{ if and only if } \mu(x) \geq p. \quad (1)$$

In other words, a p -cut of $\mu : S \rightarrow L$ is the inverse image of the principal filter $\uparrow p$, generated by $p \in L$:

$$\mu_p = \mu^{-1}(\uparrow p). \quad (2)$$

It is obvious that for every $p, q \in L$, $p \leq q$ implies $\mu_q \subseteq \mu_p$.

Cut set (p -cut)

Let $p \in L$. A **cut set** of an L -valued function $\mu : S \rightarrow L$ (a p -cut) is a subset $\mu_p \subseteq S$ defined by:

$$x \in \mu_p \text{ if and only if } \mu(x) \geq p. \quad (1)$$

In other words, a p -cut of $\mu : S \rightarrow L$ is the inverse image of the principal filter $\uparrow p$, generated by $p \in L$:

$$\mu_p = \mu^{-1}(\uparrow p). \quad (2)$$

It is obvious that for every $p, q \in L$, $p \leq q$ implies $\mu_q \subseteq \mu_p$.

Cuts and closure systems

If $\mu : S \rightarrow L$ is an L -valued function on S , then the collection μ_L of all cuts of μ is a closure system on S under the set-inclusion.

Let \mathcal{F} be a closure system on a set S . Then there is a lattice L and an L -valued function $\mu : S \rightarrow L$, such that the collection μ_L of cuts of μ is \mathcal{F} .

A required lattice L is the collection \mathcal{F} ordered by the reversed-inclusion, and that $\mu : S \rightarrow L$ can be defined as follows:

$$\mu(x) = \bigcap \{f \in \mathcal{F} \mid x \in f\}. \quad (3)$$

Cuts and closure systems

If $\mu : S \rightarrow L$ is an L -valued function on S , then the collection μ_L of all cuts of μ is a closure system on S under the set-inclusion.

Let \mathcal{F} be a closure system on a set S . Then there is a lattice L and an L -valued function $\mu : S \rightarrow L$, such that the collection μ_L of cuts of μ is \mathcal{F} .

A required lattice L is the collection \mathcal{F} ordered by the *reversed-inclusion*, and that $\mu : S \rightarrow L$ can be defined as follows:

$$\mu(x) = \bigcap \{f \in \mathcal{F} \mid x \in f\}. \quad (3)$$

Cuts and closure systems

If $\mu : S \rightarrow L$ is an L -valued function on S , then the collection μ_L of all cuts of μ is a closure system on S under the set-inclusion.

Let \mathcal{F} be a closure system on a set S . Then there is a lattice L and an L -valued function $\mu : S \rightarrow L$, such that the collection μ_L of cuts of μ is \mathcal{F} .

A required lattice L is the collection \mathcal{F} ordered by the *reversed-inclusion*, and that $\mu : S \rightarrow L$ can be defined as follows:

$$\mu(x) = \bigcap \{f \in \mathcal{F} \mid x \in f\}. \quad (3)$$

The relation \approx on L

Given an L -valued function $\mu : S \rightarrow L$, we define the relation \approx on L : for $p, q \in L$

$$p \approx q \text{ if and only if } \mu_p = \mu_q. \quad (4)$$

The relation \approx is an equivalence on L , and

$$p \approx q \text{ if and only if } \uparrow p \cap \mu(S) = \uparrow q \cap \mu(S), \quad (5)$$

where $\mu(S) = \{r \in L \mid r = \mu(x) \text{ for some } x \in S\}$.

We denote by L/\approx the collection of equivalence classes under \approx .

The relation \approx on L

Given an L -valued function $\mu : S \rightarrow L$, we define the relation \approx on L : for $p, q \in L$

$$p \approx q \text{ if and only if } \mu_p = \mu_q. \quad (4)$$

The relation \approx is an equivalence on L , and

$$p \approx q \text{ if and only if } \uparrow p \cap \mu(S) = \uparrow q \cap \mu(S), \quad (5)$$

where $\mu(S) = \{r \in L \mid r = \mu(x) \text{ for some } x \in S\}$.

We denote by L/\approx the collection of equivalence classes under \approx .

The relation \approx on L

Given an L -valued function $\mu : S \rightarrow L$, we define the relation \approx on L : for $p, q \in L$

$$p \approx q \text{ if and only if } \mu_p = \mu_q. \quad (4)$$

The relation \approx is an equivalence on L , and

$$p \approx q \text{ if and only if } \uparrow p \cap \mu(S) = \uparrow q \cap \mu(S), \quad (5)$$

where $\mu(S) = \{r \in L \mid r = \mu(x) \text{ for some } x \in S\}$.

We denote by L/\approx the collection of equivalence classes under \approx .

The collection of cuts

Let (μ_L, \leq) be the poset with $\mu_L = \{\mu_p \mid p \in L\}$ (the collection of cuts of μ) and the order \leq being the inverse of the set-inclusion: for $\mu_p, \mu_q \in \mu_L$,

$$\mu_p \leq \mu_q \text{ if and only if } \mu_q \subseteq \mu_p.$$

(μ_L, \leq) is a complete lattice and for every collection $\{\mu_p \mid p \in L_1\}$, $L_1 \subseteq L$ of cuts of μ , we have

$$\bigcap \{\mu_p \mid p \in L_1\} = \mu_{\vee(p|p \in L_1)}. \quad (6)$$

The collection of cuts

Let (μ_L, \leq) be the poset with $\mu_L = \{\mu_p \mid p \in L\}$ (the collection of cuts of μ) and the order \leq being the inverse of the set-inclusion: for $\mu_p, \mu_q \in \mu_L$,

$$\mu_p \leq \mu_q \text{ if and only if } \mu_q \subseteq \mu_p.$$

(μ_L, \leq) is a complete lattice and for every collection $\{\mu_p \mid p \in L_1\}$, $L_1 \subseteq L$ of cuts of μ , we have

$$\bigcap \{\mu_p \mid p \in L_1\} = \mu_{\vee(p|p \in L_1)}. \quad (6)$$

The quotient L/\approx

Each \approx -class contains its supremum:

$$\bigvee [p]_{\approx} \in [p]_{\approx}. \quad (7)$$

The mapping $p \mapsto \bigvee [p]_{\approx}$ is a closure operator on L .

The quotient L/\approx can be ordered by the relation $\leq_{L/\approx}$ defined as follows:

$$[p]_{\approx} \leq_{L/\approx} [q]_{\approx} \text{ if and only if } \uparrow q \cap \mu(S) \subseteq \uparrow p \cap \mu(S).$$

The order $\leq_{L/\approx}$ of classes in L/\approx corresponds to the order of suprema of classes in L (we denote the order in L by \leq_L):

The quotient L/\approx

Each \approx -class contains its supremum:

$$\bigvee [p]_{\approx} \in [p]_{\approx}. \quad (7)$$

The mapping $p \mapsto \bigvee [p]_{\approx}$ is a closure operator on L .

The quotient L/\approx can be ordered by the relation $\leq_{L/\approx}$ defined as follows:

$$[p]_{\approx} \leq_{L/\approx} [q]_{\approx} \text{ if and only if } \uparrow q \cap \mu(S) \subseteq \uparrow p \cap \mu(S).$$

The order $\leq_{L/\approx}$ of classes in L/\approx corresponds to the order of suprema of classes in L (we denote the order in L by \leq_L):

The quotient L/\approx

Each \approx -class contains its supremum:

$$\bigvee [p]_{\approx} \in [p]_{\approx}. \quad (7)$$

The mapping $p \mapsto \bigvee [p]_{\approx}$ is a closure operator on L .

The quotient L/\approx can be ordered by the relation $\leq_{L/\approx}$ defined as follows:

$$[p]_{\approx} \leq_{L/\approx} [q]_{\approx} \text{ if and only if } \uparrow q \cap \mu(S) \subseteq \uparrow p \cap \mu(S).$$

The order $\leq_{L/\approx}$ of classes in L/\approx corresponds to the order of suprema of classes in L (we denote the order in L by \leq_L):

The poset $(L/\approx, \leq_{L/\approx})$

The poset $(L/\approx, \leq_{L/\approx})$ is a complete lattice fulfilling:

- (i) $[p]_{\approx} \leq_{L/\approx} [q]_{\approx}$ if and only if $\bigvee [p]_{\approx} \leq_L \bigvee [q]_{\approx}$.*
- (ii) The mapping $[p]_{\approx} \mapsto \bigvee [p]_{\approx}$ is an injection of L/\approx into L .*

The sub-poset $(\bigvee [p]_{\approx}, \leq_L)$ of L is isomorphic to the lattice $(L/\approx, \leq_{L/\approx})$ under $\bigvee [p]_{\approx} \mapsto [p]_{\approx}$.

Let $\mu : S \rightarrow L$ be an L -valued function on S . The lattice (μ_L, \leq) of cuts of μ is isomorphic with the lattice $(L/\approx, \leq_{L/\approx})$ of \approx -classes in L under the mapping $\mu_p \mapsto [p]_{\approx}$.

The poset $(L/\approx, \leq_{L/\approx})$

The poset $(L/\approx, \leq_{L/\approx})$ is a complete lattice fulfilling:

(i) $[p]_{\approx} \leq_{L/\approx} [q]_{\approx}$ if and only if $\bigvee [p]_{\approx} \leq_L \bigvee [q]_{\approx}$.

(ii) The mapping $[p]_{\approx} \mapsto \bigvee [p]_{\approx}$ is an injection of L/\approx into L .

The sub-poset $(\bigvee [p]_{\approx}, \leq_L)$ of L is isomorphic to the lattice $(L/\approx, \leq_{L/\approx})$ under $\bigvee [p]_{\approx} \mapsto [p]_{\approx}$.

Let $\mu : S \rightarrow L$ be an L -valued function on S . The lattice (μ_L, \leq) of cuts of μ is isomorphic with the lattice $(L/\approx, \leq_{L/\approx})$ of \approx -classes in L under the mapping $\mu_p \mapsto [p]_{\approx}$.

The poset $(L/\approx, \leq_{L/\approx})$

The poset $(L/\approx, \leq_{L/\approx})$ is a complete lattice fulfilling:

- (i) $[p]_{\approx} \leq_{L/\approx} [q]_{\approx}$ if and only if $\bigvee [p]_{\approx} \leq_L \bigvee [q]_{\approx}$.*
- (ii) The mapping $[p]_{\approx} \mapsto \bigvee [p]_{\approx}$ is an injection of L/\approx into L .*

The sub-poset $(\bigvee [p]_{\approx}, \leq_L)$ of L is isomorphic to the lattice $(L/\approx, \leq_{L/\approx})$ under $\bigvee [p]_{\approx} \mapsto [p]_{\approx}$.

Let $\mu : S \rightarrow L$ be an L -valued function on S . The lattice (μ_L, \leq) of cuts of μ is isomorphic with the lattice $(L/\approx, \leq_{L/\approx})$ of \approx -classes in L under the mapping $\mu_p \mapsto [p]_{\approx}$.

The poset $(L/\approx, \leq_{L/\approx})$

The poset $(L/\approx, \leq_{L/\approx})$ is a complete lattice fulfilling:

- (i) $[p]_{\approx} \leq_{L/\approx} [q]_{\approx}$ if and only if $\bigvee [p]_{\approx} \leq_L \bigvee [q]_{\approx}$.*
- (ii) The mapping $[p]_{\approx} \mapsto \bigvee [p]_{\approx}$ is an injection of L/\approx into L .*

The sub-poset $(\bigvee [p]_{\approx}, \leq_L)$ of L is isomorphic to the lattice $(L/\approx, \leq_{L/\approx})$ under $\bigvee [p]_{\approx} \mapsto [p]_{\approx}$.

Let $\mu : S \rightarrow L$ be an L -valued function on S . The lattice (μ_L, \leq) of cuts of μ is isomorphic with the lattice $(L/\approx, \leq_{L/\approx})$ of \approx -classes in L under the mapping $\mu_p \mapsto [p]_{\approx}$.

The poset $(L/\approx, \leq_{L/\approx})$

The poset $(L/\approx, \leq_{L/\approx})$ is a complete lattice fulfilling:

- (i) $[p]_{\approx} \leq_{L/\approx} [q]_{\approx}$ if and only if $\bigvee [p]_{\approx} \leq_L \bigvee [q]_{\approx}$.*
- (ii) The mapping $[p]_{\approx} \mapsto \bigvee [p]_{\approx}$ is an injection of L/\approx into L .*

The sub-poset $(\bigvee [p]_{\approx}, \leq_L)$ of L is isomorphic to the lattice $(L/\approx, \leq_{L/\approx})$ under $\bigvee [p]_{\approx} \mapsto [p]_{\approx}$.

Let $\mu : S \rightarrow L$ be an L -valued function on S . The lattice (μ_L, \leq) of cuts of μ is isomorphic with the lattice $(L/\approx, \leq_{L/\approx})$ of \approx -classes in L under the mapping $\mu_p \mapsto [p]_{\approx}$.

Canonical representation of lattice-valued functions

We take the lattice (\mathcal{F}, \leq) , where $\mathcal{F} = \mu_L \subseteq \mathcal{P}(S)$ is the collection of cuts of μ , and the order \leq is the dual of the set inclusion.

Let $\hat{\mu} : S \rightarrow \mathcal{F}$, where

$$\hat{\mu}(x) := \bigcap \{ \mu_p \in \mu_L \mid x \in \mu_p \}. \quad (8)$$

Properties:

$\hat{\mu}$ has the same cuts as μ .

$\hat{\mu}$ has one-element classes of the equivalence relation \approx .

Every $f \in \mathcal{F}$ is equal to the corresponding cut of $\hat{\mu}$.

Canonical representation of lattice-valued functions

We take the lattice (\mathcal{F}, \leq) , where $\mathcal{F} = \mu_L \subseteq \mathcal{P}(S)$ is the collection of cuts of μ , and the order \leq is the dual of the set inclusion.

Let $\hat{\mu} : S \rightarrow \mathcal{F}$, where

$$\hat{\mu}(x) := \bigcap \{ \mu_p \in \mu_L \mid x \in \mu_p \}. \quad (8)$$

Properties:

$\hat{\mu}$ has the same cuts as μ .

$\hat{\mu}$ has one-element classes of the equivalence relation \approx .

Every $f \in \mathcal{F}$ is equal to the corresponding cut of $\hat{\mu}$.

Canonical representation of lattice-valued functions

We take the lattice (\mathcal{F}, \leq) , where $\mathcal{F} = \mu_L \subseteq \mathcal{P}(S)$ is the collection of cuts of μ , and the order \leq is the dual of the set inclusion.

Let $\hat{\mu} : S \rightarrow \mathcal{F}$, where

$$\hat{\mu}(x) := \bigcap \{ \mu_p \in \mu_L \mid x \in \mu_p \}. \quad (8)$$

Properties:

$\hat{\mu}$ has the same cuts as μ .

$\hat{\mu}$ has one-element classes of the equivalence relation \approx .

Every $f \in \mathcal{F}$ is equal to the corresponding cut of $\hat{\mu}$.

Canonical representation of lattice-valued functions

We take the lattice (\mathcal{F}, \leq) , where $\mathcal{F} = \mu_L \subseteq \mathcal{P}(S)$ is the collection of cuts of μ , and the order \leq is the dual of the set inclusion.

Let $\hat{\mu} : S \rightarrow \mathcal{F}$, where

$$\hat{\mu}(x) := \bigcap \{ \mu_p \in \mu_L \mid x \in \mu_p \}. \quad (8)$$

Properties:

$\hat{\mu}$ has the same cuts as μ .

$\hat{\mu}$ has one-element classes of the equivalence relation \approx .

Every $f \in \mathcal{F}$ is equal to the corresponding cut of $\hat{\mu}$.

Canonical representation of lattice-valued functions

We take the lattice (\mathcal{F}, \leq) , where $\mathcal{F} = \mu_L \subseteq \mathcal{P}(S)$ is the collection of cuts of μ , and the order \leq is the dual of the set inclusion.

Let $\hat{\mu} : S \rightarrow \mathcal{F}$, where

$$\hat{\mu}(x) := \bigcap \{ \mu_p \in \mu_L \mid x \in \mu_p \}. \quad (8)$$

Properties:

$\hat{\mu}$ has the same cuts as μ .

$\hat{\mu}$ has one-element classes of the equivalence relation \approx .

Every $f \in \mathcal{F}$ is equal to the corresponding cut of $\hat{\mu}$.

Canonical representation of lattice-valued functions

We take the lattice (\mathcal{F}, \leq) , where $\mathcal{F} = \mu_L \subseteq \mathcal{P}(S)$ is the collection of cuts of μ , and the order \leq is the dual of the set inclusion.

Let $\hat{\mu} : S \rightarrow \mathcal{F}$, where

$$\hat{\mu}(x) := \bigcap \{ \mu_p \in \mu_L \mid x \in \mu_p \}. \quad (8)$$

Properties:




$\hat{\mu}$ has the same cuts as μ .

$\hat{\mu}$ has one-element classes of the equivalence relation \approx .

Every $f \in \mathcal{F}$ is equal to the corresponding cut of $\hat{\mu}$.

Example

$$S = \{a, b, c, d\}$$



$$\mu = \begin{pmatrix} a & b & c & d \\ p & s & r & t \end{pmatrix}$$

$$\nu = \begin{pmatrix} a & b & c & d \\ z & w & m & v \end{pmatrix}$$

$$\hat{\mu} = \hat{\nu} = \begin{pmatrix} a & b & c & d \\ \{a\} & \{a, b\} & \{c\} & \{c, d\} \end{pmatrix}$$

Lattice-valued Boolean functions

A **Boolean function** is a mapping $f : \{0, 1\}^n \rightarrow \{0, 1\}$, $n \in \mathbb{N}$.

A **lattice-valued Boolean function** is a mapping

$$f : \{0, 1\}^n \rightarrow L,$$

where L is a complete lattice.

We also deal with **lattice-valued n -variable functions** on a finite domain $\{0, 1, \dots, k-1\}$:

$$f : \{0, 1, \dots, k-1\}^n \rightarrow L,$$

where L is a complete lattice.

We use also **p -cuts** of lattice-valued functions as characteristic functions: for $f : \{0, 1, \dots, k-1\}^n \rightarrow L$ and $p \in L$, we have

$$f_p : \{0, 1, \dots, k-1\}^n \rightarrow \{0, 1\},$$

such that $f_p(x_1, \dots, x_n) = 1$ if and only if $f(x_1, \dots, x_n) \geq p$.

Clearly, *a cut of a lattice-valued Boolean function is (as a characteristic function) a Boolean function.*

Lattice-valued Boolean functions

A **Boolean function** is a mapping $f : \{0, 1\}^n \rightarrow \{0, 1\}$, $n \in \mathbb{N}$.

A **lattice-valued Boolean function** is a mapping

$$f : \{0, 1\}^n \rightarrow L,$$

where L is a complete lattice.

We also deal with **lattice-valued n -variable functions** on a finite domain $\{0, 1, \dots, k-1\}$:

$$f : \{0, 1, \dots, k-1\}^n \rightarrow L,$$

where L is a complete lattice.

We use also **p -cuts** of lattice-valued functions as characteristic functions: for $f : \{0, 1, \dots, k-1\}^n \rightarrow L$ and $p \in L$, we have

$$f_p : \{0, 1, \dots, k-1\}^n \rightarrow \{0, 1\},$$

such that $f_p(x_1, \dots, x_n) = 1$ if and only if $f(x_1, \dots, x_n) \geq p$.

Clearly, a *cut of a lattice-valued Boolean function is (as a characteristic function) a Boolean function.*

Lattice-valued Boolean functions

A **Boolean function** is a mapping $f : \{0, 1\}^n \rightarrow \{0, 1\}$, $n \in \mathbb{N}$.

A **lattice-valued Boolean function** is a mapping

$$f : \{0, 1\}^n \rightarrow L,$$

where L is a complete lattice.

We also deal with **lattice-valued n -variable functions** on a finite domain $\{0, 1, \dots, k - 1\}$:

$$f : \{0, 1, \dots, k - 1\}^n \rightarrow L,$$

where L is a complete lattice.

We use also p -cuts of lattice-valued functions as characteristic functions: for $f : \{0, 1, \dots, k - 1\}^n \rightarrow L$ and $p \in L$, we have

$$f_p : \{0, 1, \dots, k - 1\}^n \rightarrow \{0, 1\},$$

such that $f_p(x_1, \dots, x_n) = 1$ if and only if $f(x_1, \dots, x_n) \geq p$.

Clearly, a cut of a lattice-valued Boolean function is (as a characteristic function) a Boolean function.

Lattice-valued Boolean functions

A **Boolean function** is a mapping $f : \{0, 1\}^n \rightarrow \{0, 1\}$, $n \in \mathbb{N}$.

A **lattice-valued Boolean function** is a mapping

$$f : \{0, 1\}^n \rightarrow L,$$

where L is a complete lattice.

We also deal with **lattice-valued n -variable functions** on a finite domain $\{0, 1, \dots, k - 1\}$:

$$f : \{0, 1, \dots, k - 1\}^n \rightarrow L,$$

where L is a complete lattice.

We use also **p -cuts** of lattice-valued functions as characteristic functions: for $f : \{0, 1, \dots, k - 1\}^n \rightarrow L$ and $p \in L$, we have

$$f_p : \{0, 1, \dots, k - 1\}^n \rightarrow \{0, 1\},$$

such that $f_p(x_1, \dots, x_n) = 1$ if and only if $f(x_1, \dots, x_n) \geq p$.

Clearly, a cut of a lattice-valued Boolean function is (as a characteristic function) a Boolean function.

Lattice-valued Boolean functions

A **Boolean function** is a mapping $f : \{0, 1\}^n \rightarrow \{0, 1\}$, $n \in \mathbb{N}$.

A **lattice-valued Boolean function** is a mapping

$$f : \{0, 1\}^n \rightarrow L,$$

where L is a complete lattice.

We also deal with **lattice-valued n -variable functions** on a finite domain $\{0, 1, \dots, k - 1\}$:

$$f : \{0, 1, \dots, k - 1\}^n \rightarrow L,$$

where L is a complete lattice.

We use also **p -cuts** of lattice-valued functions as characteristic functions: for $f : \{0, 1, \dots, k - 1\}^n \rightarrow L$ and $p \in L$, we have

$$f_p : \{0, 1, \dots, k - 1\}^n \rightarrow \{0, 1\},$$

such that $f_p(x_1, \dots, x_n) = 1$ if and only if $f(x_1, \dots, x_n) \geq p$.

Clearly, *a cut of a lattice-valued Boolean function is (as a characteristic function) a Boolean function.*

As usual, by S_n we denote the symmetric group of all permutations over an n -element set. If f is an n -variable function on a finite domain X and $\sigma \in S_n$, then f is **invariant** under σ , symbolically $\sigma \vdash f$, if for all $(x_1, \dots, x_n) \in X^n$

$$f(x_1, \dots, x_n) = f(x_{\sigma(1)}, \dots, x_{\sigma(n)}).$$

If f is invariant under all permutations in $G \leq S_n$ and not invariant under any permutation from $S_n \setminus G$, then G is called **the invariance group** of f , and it is denoted by $G(f)$.

As usual, by S_n we denote the symmetric group of all permutations over an n -element set. If f is an n -variable function on a finite domain X and $\sigma \in S_n$, then f is **invariant** under σ , symbolically $\sigma \vdash f$, if for all $(x_1, \dots, x_n) \in X^n$

$$f(x_1, \dots, x_n) = f(x_{\sigma(1)}, \dots, x_{\sigma(n)}).$$

If f is invariant under all permutations in $G \leq S_n$ and not invariant under any permutation from $S_n \setminus G$, then G is called **the invariance group** of f , and it is denoted by $G(f)$.

A group $G \leq S_n$ is said to be (k, m) -representable if there is a function $f : \{0, 1, \dots, k-1\}^n \rightarrow \{1, \dots, m\}$ whose invariance group is G .

If G is the invariance group of a function $f : \{0, 1, \dots, k-1\}^n \rightarrow \mathbb{N}$, then it is called (k, ∞) -representable.

$G \leq S_n$ is called m -representable if it is the invariance group of a function $f : \{0, 1\}^n \rightarrow \{1, \dots, m\}$;

it is called representable if it is m -representable for some $m \in \mathbb{N}$.

By the above, representability is equivalent with $(2, \infty)$ -representability.

A group $G \leq S_n$ is said to be (k, m) -representable if there is a function $f : \{0, 1, \dots, k-1\}^n \rightarrow \{1, \dots, m\}$ whose invariance group is G .

If G is the invariance group of a function $f : \{0, 1, \dots, k-1\}^n \rightarrow \mathbb{N}$, then it is called (k, ∞) -representable.

$G \leq S_n$ is called m -representable if it is the invariance group of a function $f : \{0, 1\}^n \rightarrow \{1, \dots, m\}$;

it is called representable if it is m -representable for some $m \in \mathbb{N}$.

By the above, representability is equivalent with $(2, \infty)$ -representability.

A group $G \leq S_n$ is said to be (k, m) -representable if there is a function $f : \{0, 1, \dots, k-1\}^n \rightarrow \{1, \dots, m\}$ whose invariance group is G .

If G is the invariance group of a function $f : \{0, 1, \dots, k-1\}^n \rightarrow \mathbb{N}$, then it is called (k, ∞) -representable.

$G \leq S_n$ is called m -representable if it is the invariance group of a function $f : \{0, 1\}^n \rightarrow \{1, \dots, m\}$;

it is called *representable* if it is m -representable for some $m \in \mathbb{N}$.

By the above, representability is equivalent with $(2, \infty)$ -representability.

A group $G \leq S_n$ is said to be (k, m) -representable if there is a function $f : \{0, 1, \dots, k-1\}^n \rightarrow \{1, \dots, m\}$ whose invariance group is G .

If G is the invariance group of a function $f : \{0, 1, \dots, k-1\}^n \rightarrow \mathbb{N}$, then it is called (k, ∞) -representable.

$G \leq S_n$ is called m -representable if it is the invariance group of a function $f : \{0, 1\}^n \rightarrow \{1, \dots, m\}$;

it is called representable if it is m -representable for some $m \in \mathbb{N}$.

By the above, representability is equivalent with $(2, \infty)$ -representability.

A group $G \leq S_n$ is said to be (k, m) -representable if there is a function $f : \{0, 1, \dots, k-1\}^n \rightarrow \{1, \dots, m\}$ whose invariance group is G .

If G is the invariance group of a function $f : \{0, 1, \dots, k-1\}^n \rightarrow \mathbb{N}$, then it is called (k, ∞) -representable.

$G \leq S_n$ is called m -representable if it is the invariance group of a function $f : \{0, 1\}^n \rightarrow \{1, \dots, m\}$;

it is called representable if it is m -representable for some $m \in \mathbb{N}$.

By the above, representability is equivalent with $(2, \infty)$ -representability.

Representability by lattice-valued functions

We say that a permutation group $G \leq S_n$ is (k, L) -**representable**, if there is a lattice-valued function $f : \{0, 1, \dots, k - 1\}^n \rightarrow L$, such that $\sigma \vdash f$ if and only if $\sigma \in G$.

In particular, a $(2, L)$ -representable group is the invariance group of a lattice-valued Boolean function $f : \{0, 1\}^n \rightarrow L$.

The notion of $(2, L)$ -representability is more general than $(2, 2)$ -representability. An example is the Klein 4-group: $\{id, (12)(34), (13)(24), (14)(23)\}$, which is $(2, L)$ representable (for L being a three element chain), but not $(2, 2)$ -representable.

Representability by lattice-valued functions

We say that a permutation group $G \leq S_n$ is (k, L) -**representable**, if there is a lattice-valued function $f : \{0, 1, \dots, k - 1\}^n \rightarrow L$, such that $\sigma \vdash f$ if and only if $\sigma \in G$.

In particular, a $(2, L)$ -representable group is the invariance group of a lattice-valued Boolean function $f : \{0, 1\}^n \rightarrow L$.

The notion of $(2, L)$ -representability is more general than $(2, 2)$ -representability. An example is the Klein 4-group: $\{id, (12)(34), (13)(24), (14)(23)\}$, which is $(2, L)$ representable (for L being a three element chain), but not $(2, 2)$ -representable.

Representability by lattice-valued functions

We say that a permutation group $G \leq S_n$ is (k, L) -**representable**, if there is a lattice-valued function $f : \{0, 1, \dots, k-1\}^n \rightarrow L$, such that $\sigma \vdash f$ if and only if $\sigma \in G$.

In particular, a $(2, L)$ -representable group is the invariance group of a lattice-valued Boolean function $f : \{0, 1\}^n \rightarrow L$.

The notion of $(2, L)$ -representability is more general than $(2, 2)$ -representability. An example is the Klein 4-group: $\{id, (12)(34), (13)(24), (14)(23)\}$, which is $(2, L)$ representable (for L being a three element chain), but not $(2, 2)$ -representable.

A Galois connection for invariance groups

Let $O_k^{(n)} = \{f \mid f: \mathbf{k}^n \rightarrow \mathbf{k}\}$ denote the set of all n -ary operations on \mathbf{k} , and for $F \subseteq O_k^{(n)}$ and $G \subseteq S_n$ let

$$F^+ := \{\sigma \in S_n \mid \forall f \in F : \sigma \vdash f\}, \quad \overline{F}^{(k)} := (F^+)^+,$$
$$G^+ := \{f \in O_k^{(n)} \mid \forall \sigma \in G : \sigma \vdash f\}, \quad \overline{G}^{(k)} := (G^+)^+.$$

The assignment $G \mapsto \overline{G}^{(k)}$ is a closure operator on S_n , and it is easy to see that $\overline{G}^{(k)}$ is a subgroup of S_n for every subset $G \subseteq S_n$ (even if G is not a group). For $G \leq S_n$, we call $\overline{G}^{(k)}$ the *Galois closure of G over \mathbf{k}* , and we say that G is *Galois closed over \mathbf{k}* if $\overline{G}^{(k)} = G$.

A Galois connection for invariance groups

Let $O_k^{(n)} = \{f \mid f: \mathbf{k}^n \rightarrow \mathbf{k}\}$ denote the set of all n -ary operations on \mathbf{k} , and for $F \subseteq O_k^{(n)}$ and $G \subseteq S_n$ let

$$F^+ := \{\sigma \in S_n \mid \forall f \in F : \sigma \vdash f\}, \quad \overline{F}^{(k)} := (F^+)^+,$$
$$G^+ := \{f \in O_k^{(n)} \mid \forall \sigma \in G : \sigma \vdash f\}, \quad \overline{G}^{(k)} := (G^+)^+.$$

The assignment $G \mapsto \overline{G}^{(k)}$ is a closure operator on S_n , and it is easy to see that $\overline{G}^{(k)}$ is a subgroup of S_n for every subset $G \subseteq S_n$ (even if G is not a group). For $G \leq S_n$, we call $\overline{G}^{(k)}$ the *Galois closure of G over \mathbf{k}* , and we say that G is *Galois closed over \mathbf{k}* if $\overline{G}^{(k)} = G$.

Galois closed groups

A group $G \leq S_n$ is Galois closed over \mathbf{k} if and only if G is (k, ∞) -representable.

For every $G \leq S_n$, we have

$$\overline{G}^{(k)} = \bigcap_{a \in \mathbf{k}^n} (S_n)_a \cdot G.$$

For arbitrary $k, n \geq 2$, characterize those subgroups of S_n that are Galois closed over \mathbf{k} .

Theorem (H., Makay, Pöschel, Waldhauser) Let $n > \max(2^d, d^2 + d)$ and $G \leq S_n$. Then G is not Galois closed over \mathbf{k} if and only if $G = A_B \times L$ or $G <_{\text{sd}} S_B \times L$, where $B \subseteq \mathbf{n}$ is such that $D := \mathbf{n} \setminus B$ has less than d elements, and L is an arbitrary permutation group on D .

Galois closed groups

A group $G \leq S_n$ is Galois closed over \mathbf{k} if and only if G is (k, ∞) -representable.

For every $G \leq S_n$, we have

$$\overline{G}^{(k)} = \bigcap_{a \in \mathbf{k}^n} (S_n)_a \cdot G.$$

For arbitrary $k, n \geq 2$, characterize those subgroups of S_n that are Galois closed over \mathbf{k} .

Theorem (H., Makay, Pöschel, Waldhauser) Let $n > \max(2^d, d^2 + d)$ and $G \leq S_n$. Then G is not Galois closed over \mathbf{k} if and only if $G = A_B \times L$ or $G <_{\text{sd}} S_B \times L$, where $B \subseteq \mathbf{n}$ is such that $D := \mathbf{n} \setminus B$ has less than d elements, and L is an arbitrary permutation group on D .

Galois closed groups

A group $G \leq S_n$ is Galois closed over \mathbf{k} if and only if G is (k, ∞) -representable.

For every $G \leq S_n$, we have

$$\overline{G}^{(k)} = \bigcap_{a \in \mathbf{k}^n} (S_n)_a \cdot G.$$

For arbitrary $k, n \geq 2$, characterize those subgroups of S_n that are Galois closed over \mathbf{k} .

Theorem (H., Makay, Pöschel, Waldhauser) Let $n > \max(2^d, d^2 + d)$ and $G \leq S_n$. Then G is not Galois closed over \mathbf{k} if and only if $G = A_B \times L$ or $G <_{\text{sd}} S_B \times L$, where $B \subseteq \mathbf{n}$ is such that $D := \mathbf{n} \setminus B$ has less than d elements, and L is an arbitrary permutation group on D .

Galois closed groups

A group $G \leq S_n$ is Galois closed over \mathbf{k} if and only if G is (k, ∞) -representable.

For every $G \leq S_n$, we have

$$\overline{G}^{(k)} = \bigcap_{a \in \mathbf{k}^n} (S_n)_a \cdot G.$$

For arbitrary $k, n \geq 2$, characterize those subgroups of S_n that are Galois closed over \mathbf{k} .

Theorem (H., Makay, Pöschel, Waldhauser) Let $n > \max(2^d, d^2 + d)$ and $G \leq S_n$. Then G is not Galois closed over \mathbf{k} if and only if $G = A_B \times L$ or $G <_{\text{sd}} S_B \times L$, where $B \subseteq \mathbf{n}$ is such that $D := \mathbf{n} \setminus B$ has less than d elements, and L is an arbitrary permutation group on D .

One can easily check that *a permutation group $G \subseteq S_n$ is L -representable if and only if it is Galois closed over $\mathbf{2}$.*

Similarly, it is easy to show that *a permutation group is (k, L) -representable if and only if it is Galois closed over the k -element domain.*

One can easily check that *a permutation group $G \subseteq S_n$ is L -representable if and only if it is Galois closed over $\mathbf{2}$.*

Similarly, it is easy to show that *a permutation group is (k, L) -representable if and only if it is Galois closed over the k -element domain.*

Cuts of composition of functions

Theorem Let L be a complete lattice, let $A \neq \emptyset$ be a set and let $\sigma : A \rightarrow A$, $\mu : A \rightarrow L$, $\psi : L \rightarrow L$. Then, for every $p \in L$,

$$(\sigma \circ \mu \circ \psi)_p = \sigma \circ \mu \circ \psi_p.$$

Corollary Let L be a complete lattice, let $A \neq \emptyset$ and let $\mu : A \rightarrow L$. Then the following holds.

(i) $\mu_p = \mu \circ (\mathcal{I}_L)_p$, where \mathcal{I}_L is the identity mapping $\mathcal{I}_L : L \rightarrow L$.

(ii) $(\sigma \circ \mu)_p = \sigma \circ \mu_p$, for $\sigma : A \rightarrow A$.

(iii) $(\mu \circ \psi)_p = \mu \circ \psi_p$, for $\psi : L \rightarrow L$.

Cuts of composition of functions

Theorem Let L be a complete lattice, let $A \neq \emptyset$ be a set and let $\sigma : A \rightarrow A$, $\mu : A \rightarrow L$, $\psi : L \rightarrow L$. Then, for every $p \in L$,

$$(\sigma \circ \mu \circ \psi)_p = \sigma \circ \mu \circ \psi_p.$$

Corollary Let L be a complete lattice, let $A \neq \emptyset$ and let $\mu : A \rightarrow L$. Then the following holds.

- (i) $\mu_p = \mu \circ (\mathcal{I}_L)_p$, where \mathcal{I}_L is the identity mapping $\mathcal{I}_L : L \rightarrow L$.
- (ii) $(\sigma \circ \mu)_p = \sigma \circ \mu_p$, for $\sigma : A \rightarrow A$.
- (iii) $(\mu \circ \psi)_p = \mu \circ \psi_p$, where ψ is a map $\psi : L \rightarrow L$.

Cuts of composition of functions

Theorem Let L be a complete lattice, let $A \neq \emptyset$ be a set and let $\sigma : A \rightarrow A$, $\mu : A \rightarrow L$, $\psi : L \rightarrow L$. Then, for every $p \in L$,

$$(\sigma \circ \mu \circ \psi)_p = \sigma \circ \mu \circ \psi_p.$$

Corollary Let L be a complete lattice, let $A \neq \emptyset$ and let $\mu : A \rightarrow L$. Then the following holds.

- (i) $\mu_p = \mu \circ (\mathcal{I}_L)_p$, where \mathcal{I}_L is the identity mapping $\mathcal{I}_L : L \rightarrow L$.
- (ii) $(\sigma \circ \mu)_p = \sigma \circ \mu_p$, for $\sigma : A \rightarrow A$.
- (iii) $(\mu \circ \psi)_p = \mu \circ \psi_p$, where ψ is a map $\psi : L \rightarrow L$.

Cuts of composition of functions

Theorem Let L be a complete lattice, let $A \neq \emptyset$ be a set and let $\sigma : A \rightarrow A$, $\mu : A \rightarrow L$, $\psi : L \rightarrow L$. Then, for every $p \in L$,

$$(\sigma \circ \mu \circ \psi)_p = \sigma \circ \mu \circ \psi_p.$$

Corollary Let L be a complete lattice, let $A \neq \emptyset$ and let $\mu : A \rightarrow L$. Then the following holds.

- (i) $\mu_p = \mu \circ (\mathcal{I}_L)_p$, where \mathcal{I}_L is the identity mapping $\mathcal{I}_L : L \rightarrow L$.
- (ii) $(\sigma \circ \mu)_p = \sigma \circ \mu_p$, for $\sigma : A \rightarrow A$.
- (iii) $(\mu \circ \psi)_p = \mu \circ \psi_p$, where ψ is a map $\psi : L \rightarrow L$.

Cuts of composition of functions

Theorem Let L be a complete lattice, let $A \neq \emptyset$ be a set and let $\sigma : A \rightarrow A$, $\mu : A \rightarrow L$, $\psi : L \rightarrow L$. Then, for every $p \in L$,

$$(\sigma \circ \mu \circ \psi)_p = \sigma \circ \mu \circ \psi_p.$$

Corollary Let L be a complete lattice, let $A \neq \emptyset$ and let $\mu : A \rightarrow L$. Then the following holds.

- (i) $\mu_p = \mu \circ (\mathcal{I}_L)_p$, where \mathcal{I}_L is the identity mapping $\mathcal{I}_L : L \rightarrow L$.
- (ii) $(\sigma \circ \mu)_p = \sigma \circ \mu_p$, for $\sigma : A \rightarrow A$.
- (iii) $(\mu \circ \psi)_p = \mu \circ \psi_p$, where ψ is a map $\psi : L \rightarrow L$.

Proposition Let $f : \{0, \dots, k-1\}^n \rightarrow L$ and $\sigma \in S_n$. Then

$\sigma \vdash f$ if and only if for every $p \in L$, $\sigma \vdash f_p$.

The invariance group of a lattice-valued function f depends only on the canonical representation of f .

If $f_1 : \{0, \dots, k-1\}^n \rightarrow L_1$ and $f_2 : \{0, \dots, k-1\}^n \rightarrow L_2$ are two n -variable lattice-valued functions on the same domain, then $\widehat{f}_1 = \widehat{f}_2$ implies $G(f_1) = G(f_2)$.

Proposition Let $f : \{0, \dots, k-1\}^n \rightarrow L$ and $\sigma \in S_n$. Then

$\sigma \vdash f$ if and only if for every $p \in L$, $\sigma \vdash f_p$.

The invariance group of a lattice-valued function f depends only on the canonical representation of f .

If $f_1 : \{0, \dots, k-1\}^n \rightarrow L_1$ and $f_2 : \{0, \dots, k-1\}^n \rightarrow L_2$ are two n -variable lattice-valued functions on the same domain, then $\widehat{f}_1 = \widehat{f}_2$ implies $G(f_1) = G(f_2)$.

Proposition Let $f : \{0, \dots, k-1\}^n \rightarrow L$ and $\sigma \in S_n$. Then

$\sigma \vdash f$ if and only if for every $p \in L$, $\sigma \vdash f_p$.

The invariance group of a lattice-valued function f depends only on the canonical representation of f .

If $f_1 : \{0, \dots, k-1\}^n \rightarrow L_1$ and $f_2 : \{0, \dots, k-1\}^n \rightarrow L_2$ are two n -variable lattice-valued functions on the same domain, then $\widehat{f_1} = \widehat{f_2}$ implies $G(f_1) = G(f_2)$.

Representation theorem

For every $n \in \mathbb{N}$, there is a lattice L and a lattice valued Boolean function $F : \{0, 1\}^n \rightarrow L$ satisfying the following: If $G \leq S_n$ and $G = G(f)$ for a Boolean function f , then $G = G(F_p)$, for a cut F_p .

Representation theorem on the k -element set

Every subgroups of S_n is an invariance group of a function $\{0, \dots, k-1\}^n \rightarrow \{0, \dots, k-1\}$ if and only if $k \geq n$.

If $k \geq n$, then for every subgroup G of S_n there exists a function $f : \{0, \dots, k-1\}^n \rightarrow \{0, 1\}$ such that the invariance group of f is exactly G .

For $k, n \in \mathbb{N}$ and $k \geq n$, there is a lattice L and a lattice valued function $F : \{0, \dots, k-1\}^n \rightarrow L$ such that the following holds: If $G \leq S_n$, then $G = G(F_p)$ for a cut F_p of F .

Representation theorem on the k -element set

Every subgroups of S_n is an invariance group of a function $\{0, \dots, k-1\}^n \rightarrow \{0, \dots, k-1\}$ if and only if $k \geq n$.

If $k \geq n$, then for every subgroup G of S_n there exists a function $f : \{0, \dots, k-1\}^n \rightarrow \{0, 1\}$ such that the invariance group of f is exactly G .

For $k, n \in \mathbb{N}$ and $k \geq n$, there is a lattice L and a lattice valued function $F : \{0, \dots, k-1\}^n \rightarrow L$ such that the following holds: If $G \leq S_n$, then $G = G(F_p)$ for a cut F_p of F .

Representation theorem on the k -element set

Every subgroups of S_n is an invariance group of a function $\{0, \dots, k-1\}^n \rightarrow \{0, \dots, k-1\}$ if and only if $k \geq n$.

If $k \geq n$, then for every subgroup G of S_n there exists a function $f : \{0, \dots, k-1\}^n \rightarrow \{0, 1\}$ such that the invariance group of f is exactly G .

For $k, n \in \mathbb{N}$ and $k \geq n$, there is a lattice L and a lattice valued function $F : \{0, \dots, k-1\}^n \rightarrow L$ such that the following holds: If $G \leq S_n$, then $G = G(F_p)$ for a cut F_p of F .

A *lattice-valued Boolean function* is a map $\mu: \{0, 1\}^n \rightarrow L$ where L is a bounded lattice and $n \in \langle 1, 2, 3, \dots \rangle$.

We say that μ can be given by a *linear combination* (in L) if there are $w_1, \dots, w_n \in L$ such that, for all $x = \{x_1, \dots, x_n\} \in \{0, 1\}^n$,

$$\mu(x) = \bigvee_{i=1}^n w_i x_i, \quad \text{that is,} \quad \mu(x) = \bigvee_{i=1}^n (w_i \wedge x_i). \quad (9)$$

A *lattice-valued Boolean function* is a map $\mu: \{0, 1\}^n \rightarrow L$ where L is a bounded lattice and $n \in \langle 1, 2, 3, \dots \rangle$.

We say that μ can be given by a *linear combination* (in L) if there are $w_1, \dots, w_n \in L$ such that, for all $x = \{x_1, \dots, x_n\} \in \{0, 1\}^n$,

$$\mu(x) = \bigvee_{i=1}^n w_i x_i, \quad \text{that is,} \quad \mu(x) = \bigvee_{i=1}^n (w_i \wedge x_i). \quad (9)$$

Cuts and closure systems

For $p \in L$, the set

$$\mu_p := \{x \in \{0, 1\}^n : \mu(x) \geq p\} \quad (10)$$

is called a *cut* of μ .

A *closure system* \mathcal{F} over B_n is a \cap -subsemilattice of the powerset lattice $P(B_n) = \langle P(B_n); \cup, \cap \rangle$ such that $B_n \in \mathcal{F}$. By finiteness, \mathcal{F} is necessarily a complete \cap -semilattice.

A closure system \mathcal{F} determines a *closure operator* in the standard way. We only need the closures of singleton sets, that is,

$$\text{for } x \in B_n, \text{ we have } \bar{x} := \bigcap \{f \in \mathcal{F} : x \in f\}. \quad (11)$$

For $p \in L$, the set

$$\mu_p := \{x \in \{0, 1\}^n : \mu(x) \geq p\} \quad (10)$$

is called a *cut* of μ .

A *closure system* \mathcal{F} over B_n is a \cap -subsemilattice of the powerset lattice $P(B_n) = \langle P(B_n); \cup, \cap \rangle$ such that $B_n \in \mathcal{F}$. By finiteness, \mathcal{F} is necessarily a complete \cap -semilattice.

A closure system \mathcal{F} determines a *closure operator* in the standard way. We only need the closures of singleton sets, that is,

$$\text{for } x \in B_n, \text{ we have } \bar{x} := \bigcap \{f \in \mathcal{F} : x \in f\}. \quad (11)$$

For $p \in L$, the set

$$\mu_p := \{x \in \{0, 1\}^n : \mu(x) \geq p\} \quad (10)$$

is called a *cut* of μ .

A *closure system* \mathcal{F} over B_n is a \cap -subsemilattice of the powerset lattice $P(B_n) = \langle P(B_n); \cup, \cap \rangle$ such that $B_n \in \mathcal{F}$. By finiteness, \mathcal{F} is necessarily a complete \cap -semilattice.

A closure system \mathcal{F} determines a *closure operator* in the standard way. We only need the closures of singleton sets, that is,

$$\text{for } x \in B_n, \text{ we have } \bar{x} := \bigcap \{f \in \mathcal{F} : x \in f\}. \quad (11)$$

$\{\vee, 0\}$ -homomorphism

If $\mu: B_n \rightarrow L$ such that $\mu(0) = 0$ and, for all $x, y \in B_n$,
 $\mu(x \vee y) = \mu(x) \vee \mu(y)$, then μ is called a $\{\vee, 0\}$ -homomorphism.

A lattice-valued function $B_n \rightarrow L$ can be given by a linear combination in L iff it is a $\{\vee, 0\}$ -homomorphism.

$$\mu(x \vee y) = \bigvee_i w_i (x_i \vee y_i) = \bigvee_i (w_i x_i \vee w_i y_i) = \bigvee_i w_i x_i \vee \bigvee_i w_i y_i = \mu(x) \vee \mu(y).$$

Let $e^{(i)} = \langle 0, \dots, 0, 1, 0, \dots, 0 \rangle \in B_n$ where 1 stands in the i -th place. Define $w_i := \mu(e^{(i)})$. Observe that $\mu(e^{(i)} \cdot 1) = w_i = w_i \cdot 1$ and $\mu(e^{(i)} \cdot 0) = 0 = w_i \cdot 0$, that is, $\mu(e^{(i)} \cdot x_i) = w_i \cdot x_i$. Therefore, for $x \in B_n$, we obtain $\mu(x) = \mu(\bigvee_i e^{(i)} x_i) = \bigvee_i \mu(e^{(i)} x_i) = \bigvee_i w_i \cdot x_i$.

$\{\vee, 0\}$ -homomorphism

If $\mu: B_n \rightarrow L$ such that $\mu(0) = 0$ and, for all $x, y \in B_n$, $\mu(x \vee y) = \mu(x) \vee \mu(y)$, then μ is called a $\{\vee, 0\}$ -homomorphism.

A lattice-valued function $B_n \rightarrow L$ can be given by a linear combination in L iff it is a $\{\vee, 0\}$ -homomorphism.

$$\mu(x \vee y) = \bigvee_i w_i(x_i \vee y_i) = \bigvee_i (w_i x_i \vee w_i y_i) = \bigvee_i w_i x_i \vee \bigvee_i w_i y_i = \mu(x) \vee \mu(y).$$

Let $e^{(i)} = \langle 0, \dots, 0, 1, 0, \dots, 0 \rangle \in B_n$ where 1 stands in the i -th place. Define $w_i := \mu(e^{(i)})$. Observe that $\mu(e^{(i)} \cdot 1) = w_i = w_i \cdot 1$ and $\mu(e^{(i)} \cdot 0) = 0 = w_i \cdot 0$, that is, $\mu(e^{(i)} \cdot x_i) = w_i \cdot x_i$. Therefore, for $x \in B_n$, we obtain $\mu(x) = \mu(\bigvee_i e^{(i)} x_i) = \bigvee_i \mu(e^{(i)} x_i) = \bigvee_i w_i \cdot x_i$.

$\{\vee, 0\}$ -homomorphism

If $\mu: B_n \rightarrow L$ such that $\mu(0) = 0$ and, for all $x, y \in B_n$,
 $\mu(x \vee y) = \mu(x) \vee \mu(y)$, then μ is called a $\{\vee, 0\}$ -homomorphism.

A lattice-valued function $B_n \rightarrow L$ can be given by a linear combination in L iff it is a $\{\vee, 0\}$ -homomorphism.

$$\mu(x \vee y) = \bigvee_i w_i (x_i \vee y_i) = \bigvee_i (w_i x_i \vee w_i y_i) = \bigvee_i w_i x_i \vee \bigvee_i w_i y_i = \mu(x) \vee \mu(y).$$

Let $e^{(i)} = \langle 0, \dots, 0, 1, 0, \dots, 0 \rangle \in B_n$ where 1 stands in the i -th place. Define $w_i := \mu(e^{(i)})$. Observe that $\mu(e^{(i)} \cdot 1) = w_i = w_i \cdot 1$ and $\mu(e^{(i)} \cdot 0) = 0 = w_i \cdot 0$, that is, $\mu(e^{(i)} \cdot x_i) = w_i \cdot x_i$. Therefore, for $x \in B_n$, we obtain $\mu(x) = \mu(\bigvee_i e^{(i)} x_i) = \bigvee_i \mu(e^{(i)} x_i) = \bigvee_i w_i \cdot x_i$.

$\{\vee, 0\}$ -homomorphism

If $\mu: B_n \rightarrow L$ such that $\mu(0) = 0$ and, for all $x, y \in B_n$,
 $\mu(x \vee y) = \mu(x) \vee \mu(y)$, then μ is called a $\{\vee, 0\}$ -homomorphism.

A lattice-valued function $B_n \rightarrow L$ can be given by a linear combination in L iff it is a $\{\vee, 0\}$ -homomorphism.

$$\mu(x \vee y) = \bigvee_i w_i(x_i \vee y_i) = \bigvee_i (w_i x_i \vee w_i y_i) = \bigvee_i w_i x_i \vee \bigvee_i w_i y_i = \mu(x) \vee \mu(y).$$

Let $e^{(i)} = \langle 0, \dots, 0, 1, 0, \dots, 0 \rangle \in B_n$ where 1 stands in the i -th place. Define $w_i := \mu(e^{(i)})$. Observe that $\mu(e^{(i)} \cdot 1) = w_i = w_i \cdot 1$ and $\mu(e^{(i)} \cdot 0) = 0 = w_i \cdot 0$, that is, $\mu(e^{(i)} \cdot x_i) = w_i \cdot x_i$. Therefore, for $x \in B_n$, we obtain $\mu(x) = \mu(\bigvee_i e^{(i)} x_i) = \bigvee_i \mu(e^{(i)} x_i) = \bigvee_i w_i \cdot x_i$.

If $\emptyset \neq X \subseteq B_n$ such that $(\forall x \in X)(\forall y \in B_n)(x \leq y \implies y \in X)$, then X is an *up-set* of B_n .

The lattice-valued function $\mu: B_n \rightarrow L$ is isotone iff all the cuts of μ are up-sets.

If $\emptyset \neq X \subseteq B_n$ such that $(\forall x \in X)(\forall y \in B_n)(x \leq y \text{ then } y \in X)$, then X is an *up-set* of B_n .

The lattice-valued function $\mu: B_n \rightarrow L$ is isotone iff all the cuts of μ are up-sets.

Closure systems of up-sets, linear combinations

Let \mathcal{F} be a set consisting of some up-sets of B_n . Then, the following three conditions are equivalent.

(i) \mathcal{F} is a closure system over B_n , and for all $x, y \in B_n$, $\bar{x} \subseteq \bar{y}$ implies $\overline{x \vee y} = \bar{x}$.

(ii) \mathcal{F} is a closure system over B_n , and for all $x, y \in B_n$, $\overline{x \vee y} = \bar{x} \cap \bar{y}$.

(iii) There exist a bounded lattice L and a lattice-valued function $\mu: B_n \rightarrow L$ given by a linear combination such that \mathcal{F} is the family of cuts of μ .

Closure systems of up-sets, linear combinations

Let \mathcal{F} be a set consisting of some up-sets of B_n . Then, the following three conditions are equivalent.

(i) \mathcal{F} is a closure system over B_n , and for all $x, y \in B_n$, $\bar{x} \subseteq \bar{y}$ implies $\overline{x \vee y} = \bar{x}$.

(ii) \mathcal{F} is a closure system over B_n , and for all $x, y \in B_n$, $\overline{x \vee y} = \bar{x} \cap \bar{y}$.

(iii) There exist a bounded lattice L and a lattice-valued function $\mu: B_n \rightarrow L$ given by a linear combination such that \mathcal{F} is the family of cuts of μ .

Closure systems of up-sets, linear combinations

Let \mathcal{F} be a set consisting of some up-sets of B_n . Then, the following three conditions are equivalent.

(i) \mathcal{F} is a closure system over B_n , and for all $x, y \in B_n$, $\bar{x} \subseteq \bar{y}$ implies $\overline{x \vee y} = \bar{x}$.

(ii) \mathcal{F} is a closure system over B_n , and for all $x, y \in B_n$, $\overline{x \vee y} = \bar{x} \cap \bar{y}$.

(iii) There exist a bounded lattice L and a lattice-valued function $\mu: B_n \rightarrow L$ given by a linear combination such that \mathcal{F} is the family of cuts of μ .

Let \mathcal{F} be a set consisting of some up-sets of B_n . Then, the following three conditions are equivalent.

(i) \mathcal{F} is a closure system over B_n , and for all $x, y \in B_n$, $\bar{x} \subseteq \bar{y}$ implies $\overline{x \vee y} = \bar{x}$.

(ii) \mathcal{F} is a closure system over B_n , and for all $x, y \in B_n$, $\overline{x \vee y} = \bar{x} \cap \bar{y}$.

(iii) There exist a bounded lattice L and a lattice-valued function $\mu: B_n \rightarrow L$ given by a linear combination such that \mathcal{F} is the family of cuts of μ .

Thank you for your attention!

Thank you for your attention!