

Finslerian hypersurfaces of a Finsler space with deformed Randers (α, β) - metric

V. K. Chaubey¹ and Brijesh Kumar Tripathi^{2 3}

Abstract. In the present paper, we have studied the Finslerian hypersurfaces of a Finsler Space with deformed Randers (α, β) -metric. Further, we have endeavored to prove the conditions under which these hypersurfaces of a special Finsler space with deformed Randers (α, β) metric will be a hyperplane of the first, second and third kinds.

AMS Mathematics Subject Classification (2010): 53B40 ; 53C60

Key words and phrases: Finslerian hypersurface; nonholonomic frames; Randers metric; Riemannian metric and one form metric

1. Introduction

Finsler metric $L(\alpha, \beta)$ of two variables is known as (α, β) -metric, where L is positively homogeneous function of degree one in an n -dimensional differentiable manifold M^n and the space $F^n = \{M^n, L\}$ is Finsler space with (α, β) -metric. The interesting and important examples of an (α, β) -metric are Randers metric $\alpha + \beta$, Kropina metric $\frac{\alpha^2}{\beta}$ and Matsumoto metric $\frac{\alpha^2}{(\alpha - \beta)}$ [7, 9]. Matsumoto [7] studied these metrics in detail and obtained various interesting geometrical properties which shows the importance of these metrics in the development of Finsler geometry.

In 2018, Tripathi and Chaubey [12] considered two different deformed special (α, β) -metrics of degree two in which one was formed as the combination of Randers and Riemannian metric and determined the nonholonomic frame due to this metric. The nonholonomic frame is to be considering if the metric defined by:

Definition 1.1. A space $F^n = \{M^n, F(x, y)\}$ is said to be Finsler space with (α, β) -metric, if there exists a 2-homogeneous function L of two variables such that the Finsler metric $F : TM \rightarrow \mathbb{R}$ is given by

$$F^2(x, y) = L\{\alpha(x, y), \beta(x, y)\},$$

where $\alpha^2(x, y) = a_{ij}(x)y^i y^j$, α is a Riemannian metric on the manifold M^n and $\beta(x, y) = b_i(x)y^i$ is a 1-form on the manifold M^n .

¹Department of Mathematics, North Eastern Hill University, Shillong-793022, Meghalaya, India,
e-mail: vkchaubey@outlook.com

²Department of Mathematics, L. D. College of Engineering, Navrangpura, Ahmedabad (Gujarat)-380015, India,
e-mail: brijeshkumartripathi4@gmail.com

³Corresponding author

The special (α, β) -metric [12] which is expressed in the form

$$(1.1) \quad L = (\alpha + \beta)\alpha = \alpha^2 + \alpha\beta,$$

where $\alpha = \sqrt{a_{ij}(x)y^i y^j}$ is a Riemannian metric and $\beta = b_i(x)y^i$ is a 1-form with $\|\beta_x\|_\alpha < b_0$, $\forall x \in M^n$ is known as deformed Randers (α, β) -metric.

The concept of Finslerian hypersurface was first introduced by Matsumoto [6] in 1985 and where he obtained the three types of hypersurfaces that were called hyperplane of the first, second and third kinds. Moreover, many authors [1, 2, 3, 4, 5, 8, 10, 11] studied the Finslerian hypersurfaces for the Finsler space equipped with 1-degree homogeneous Finsler metric and obtained the conditions under which it will be a hyperplane of the first, second and third kinds.

In the present paper, we have considered the deformed Randers (α, β) -metric which forms a nonholonomic frame of (α, β) -metric [12] and a 2-degree homogeneous function written in equation (1.1). Further, we have examined the conditions under these hypersurfaces of a special Finsler space with deformed Randers (α, β) metric will be a hyperplane of the first, second and third kinds.

2. Finsler spaces F^n with deformed Randers (α, β) -metric

In the present paper we consider an n -dimensional Finsler space $F^n = \{M^n, L(\alpha, \beta)\}$, that is equipped with deformed Randers (α, β) -metric which is given by the equation (1.1).

Differentiating equation (1.1) partially with respect to α and β , we have

$$L_\alpha = 2\alpha + \beta, \quad L_\beta = \alpha, \quad L_{\alpha\alpha} = 2,$$

$$L_{\beta\beta} = 0 \quad \text{and} \quad L_{\alpha\beta} = 1,$$

$$\text{where } L_\alpha = \frac{\partial L}{\partial \alpha}, \quad L_\beta = \frac{\partial L}{\partial \beta}, \quad L_{\alpha\alpha} = \frac{\partial L_\alpha}{\partial \alpha}, \quad L_{\beta\beta} = \frac{\partial L_\beta}{\partial \beta}, \quad L_{\alpha\beta} = \frac{\partial L_\alpha}{\partial \beta}.$$

In the Finsler space $F^n = \{M^n, L(\alpha, \beta)\}$ the normalized element of support $l_i = \partial_i L$ and angular metric tensor h_{ij} are given by [7]

$$(2.1) \quad l_i = \alpha^{-1} L_\alpha Y_i + L_\beta b_i,$$

and

$$(2.2) \quad h_{ij} = p a_{ij} + q_0 b_i b_j + q_{-1} (b_i Y_j + b_j Y_i) + q_{-2} Y_i Y_j,$$

where $Y_i = a_{ij} y^j$. For the fundamental metric (1.1) the above constants are

$$(2.3) \quad p = 2\alpha^2, \quad q_0 = 0, \quad q_{-1} = \alpha, \quad q_{-2} = 0.$$

The fundamental metric tensor $g_{ij} = \frac{1}{2}\dot{\partial}_i\dot{\partial}_j L^2$ and its reciprocal tensor g^{ij} for $L = L(\alpha, \beta)$ are given by

$$(2.4) \quad g_{ij} = pa_{ij} + p_0b_ib_j + p_{-1}(b_iY_j + b_jY_i) + p_{-2}Y_iY_j,$$

where

$$(2.5) \quad \begin{cases} p_0 = q_0 + L_\beta^2 = \alpha^2, \\ p_{-1} = q_{-1} + L^{-1}pL_\beta = 3\alpha, \\ p_{-2} = q_{-2} + p^2L^{-2} = 4. \end{cases}$$

The reciprocal tensor g^{ij} of g_{ij} is given by

$$(2.6) \quad g^{ij} = p^{-1}a^{ij} - s_0b^ib^j - s_{-1}(b^iy^j + b^jy^i) - s_{-2}y^iy^j,$$

where $b^i = a^{ij}b_j$, $b^2 = a_{ij}b^ib^j$ and

$$(2.7) \quad \begin{cases} s_0 = \frac{1}{\tau p}\{pp_0 + (p_0p_{-2} - p_{-1}^2)\alpha^2\}, \\ s_{-1} = \frac{1}{\tau p}\{pp_{-1} + (p_0p_{-2} - p_{-1}^2)\beta\}, \\ s_{-2} = \frac{1}{\tau p}\{pp_{-2} + (p_0p_{-2} - p_{-1}^2)b^2\}, \\ \tau = p(p + p_0b^2 + p_{-1}\beta) + (p_0p_{-2} - p_{-1}^2)(\alpha^2b^2 - \beta^2). \end{cases}$$

The hv-torsion tensor $C_{ijk} = \frac{1}{2}\dot{\partial}_k g_{ij}$ is given by

$$(2.8) \quad 2pC_{ijk} = p_{-1}(h_{ij}m_k + h_{jk}m_i + h_{ki}m_j) + \gamma_1m_im_jm_k,$$

where

$$(2.9) \quad \gamma_1 = p\frac{\partial p_0}{\partial \beta} - 3p_{-1}q_0, \quad m_i = b_i - \alpha^{-2}\beta Y_i.$$

Here m_i is a non-vanishing covariant vector orthogonal to the element of support y^i . Thus we have

Proposition 2.1. *The normalised supporting element l_i and angular metric tensor h_{ij} of an n -dimensional Finsler space F^n equipped with a deformed Randers (α, β) metric L are given by (2.1) and (2.2), respectively.*

Proposition 2.2. *The fundamental metric tensor g_{ij} and its reciprocal tensor g^{ij} of an n -dimensional Finsler space F^n equipped with a deformed Randers (α, β) metric L are given by (2.4) and (2.6), respectively.*

Proposition 2.3. *The Cartan hv-torsion tensor of an n -dimensional Finsler space F^n equipped with a deformed Randers (α, β) metric L is given by (2.8).*

Let $\{^i_{jk}\}$ be the component of Christoffel symbols for the associated Riemannian space R^n and ∇_k be the covariant derivative with respect to x^k relative to Christoffel symbol. Now we define

$$(2.10) \quad 2E_{ij} = b_{ij} + b_{ji}, \quad 2F_{ij} = b_{ij} - b_{ji},$$

where $b_{ij} = \nabla_j b_i$.

Let $CT = (\Gamma_{jk}^{*i}, \Gamma_{0k}^{*i}, \Gamma_{jk}^i)$ be the Cartan connection of F^n . The difference tensor $D_{jk}^i = \Gamma_{jk}^{*i} - \{\}_{jk}^i$ of the special Finsler space F^n is given by

$$(2.11) \quad \left\{ \begin{array}{l} D_{jk}^i = B^i E_{jk} + F_k^i B_j + F_j^i B_k + B_j^i b_{0k} + B_k^i b_{0j} \\ \quad - b_{0m} g^{im} B_{jk} - C_{jm}^i A_k^m - C_{km}^i A_j^m + C_{jkm} A_s^m g^{is} \\ \quad + \lambda^s (C_{jm}^i C_{sk}^m + C_{km}^i C_{sj}^m - C_{jk}^m C_{ms}^i), \end{array} \right.$$

where

$$(2.12) \quad \left\{ \begin{array}{l} B_k = p_0 b_k + p_{-1} Y_k, \\ B^i = g^{ij} B_j, \\ F_i^k = g^{kj} F_{ji}, \\ B_{ij} = \frac{\{p_{-1}(a_{ij} - \alpha^{-2} Y_i Y_j) + \frac{\partial p_0}{\partial \beta} m_i m_j\}}{2}, \\ B_i^k = g^{kj} B_{ji}, \\ A_k^m = B_k^m E_{00} + B^m E_{k0} + B_k F_0^m + B_0 F_k^m, \\ \lambda^m = B^m E_{00} + 2B_0 F_0^m, \\ B_0 = B_i y^i. \end{array} \right.$$

Here ' ∂ ' denotes contraction with y^i except the quantities p_0, q_0 and s_o . Thus, we have

Proposition 2.4. *The difference tensor D_{jk}^i of the Cartan connection CT for the n -dimensional Finsler space F^n is equipped with a deformed Randers (α, β) metric L is given by (2.12).*

3. Induced Cartan connection

Let $F^{(n-1)}$ be a hypersurface of F^n given by the equation $x^i = x^i(u^\alpha)$ where $\{\alpha = 1, 2, 3, \dots, (n-1)\}$. The element of support y^i of F^n is to be taken as tangential to $F^{(n-1)}$, that is [6],

$$(3.1) \quad y^i = B_\alpha^i(u) v^\alpha.$$

The metric tensor $g_{\alpha\beta}$ and hv-tensor $C_{\alpha\beta\gamma}$ of $F^{(n-1)}$ are given by

$$g_{\alpha\beta} = g_{ij} B_\alpha^i B_\beta^j, \quad C_{\alpha\beta\gamma} = C_{ijk} B_\alpha^i B_\beta^j B_\gamma^k,$$

and at each point (u^α) of $F^{(n-1)}$, a unit normal vector $N^i(u, v)$ is defined by

$$g_{ij} \{x(u, v), y(u, v)\} B_\alpha^i N^j = 0, \quad g_{ij} \{x(u, v), y(u, v)\} N^i N^j = 1.$$

The angular metric tensor $h_{\alpha\beta}$ of the hypersurface is given by

$$(3.2) \quad h_{\alpha\beta} = h_{ij} B_\alpha^i B_\beta^j, \quad h_{ij} B_\alpha^i N^j = 0, \quad h_{ij} N^i N^j = 1.$$

The relations between the components of (B_i^α, N_i) and its inverse (B_α^i, N^i) is given by

$$B_i^\alpha = g^{\alpha\beta} g_{ij} B_\beta^j, \quad B_\alpha^i B_i^\beta = \delta_\alpha^\beta, \quad B_i^\alpha N^i = 0, \quad B_\alpha^i N_i = 0.$$

$$N_i = g_{ij} N^j, \quad B_i^k = g^{kj} B_{ji}, \quad B_\alpha^i B_j^\alpha + N^i N_j = \delta_j^i.$$

The induced Cartan connection $ICT = (\Gamma_{\beta\gamma}^{\alpha}, G_\beta^\alpha, C_{\beta\gamma}^\alpha)$ of $F^{(n-1)}$ from the Cartan's connection $CT = (\Gamma_{jk}^{*i}, \Gamma_{0k}^{*i}, C_{jk}^{*i})$ is given by [6],

$$\Gamma_{\beta\gamma}^{*\alpha} = B_i^\alpha (B_{\beta\gamma}^i + \Gamma_{jk}^{*i} B_\beta^j B_\gamma^k) + M_\beta^\alpha H_\gamma,$$

$$G_\beta^\alpha = B_i^\alpha (B_{0\beta}^i + \Gamma_{0j}^{*i} B_\beta^j), \quad C_{\beta\gamma}^\alpha = B_i^\alpha C_{jk}^i B_\beta^j B_\gamma^k,$$

where

$$M_{\beta\gamma} = N_i C_{jk}^i B_\beta^j B_\gamma^k, \quad M_\beta^\alpha = g^{\alpha\gamma} M_{\beta\gamma}, \quad H_\beta = N_i (B_{0\beta}^i + \Gamma_{0j}^{*i} B_\beta^j) \quad \text{and}$$

$$B_{\beta\gamma}^i = \frac{\partial B_\beta^i}{\partial u^\gamma}, \quad B_{0\beta}^i = B_{\alpha\beta}^i v^\alpha.$$

The quantities $M_{\beta\gamma}$ and H_β are called the second fundamental v-tensor and normal curvature vector, respectively [6]. The second fundamental h-tensor $H_{\beta\gamma}$ is defined as [6]

$$(3.3) \quad H_{\beta\gamma} = N_i (B_{\beta\gamma}^i + \Gamma_{jk}^{*i} B_\beta^j B_\gamma^k) + M_\beta H_\gamma,$$

where

$$(3.4) \quad M_\beta = N_i C_{jk}^i B_\beta^j N^k.$$

The relative h and v-covariant derivatives of projection factor B_α^i with respect to ICT are given by

$$B_{\alpha|\beta}^i = H_{\alpha\beta} N^i, \quad B_{\alpha}^i|_\beta = M_{\alpha\beta} N^i.$$

It is obvious from the equation (3.3) that $H_{\beta\gamma}$ is generally not symmetric and

$$(3.5) \quad H_{\beta\gamma} - H_{\gamma\beta} = M_\beta H_\gamma - M_\gamma H_\beta.$$

From above equation, we get

$$(3.6) \quad H_{0\gamma} = H_\gamma, \quad H_{\gamma 0} = H_\gamma + M_\gamma H_0.$$

We shall use the following lemmas which are due to Matsumoto [6] in the next section

Lemma 3.1. *The normal curvature $H_0 = H_\beta v^\beta$ vanishes if and only if the normal curvature vector H_β vanishes.*

Lemma 3.2. *A hypersurface $F^{(n-1)}$ is a hyperplane of the first kind with respect to connection CT if and only if $H_\alpha = 0$.*

Lemma 3.3. *A hypersurface $F^{(n-1)}$ is a hyperplane of the second kind with respect to connection CT if and only if $H_\alpha = 0$ and $H_{\alpha\beta} = 0$.*

Lemma 3.4. *A hypersurface $F^{(n-1)}$ is a hyperplane of the third kind with respect to connection CT if and only if $H_\alpha = 0$ and $H_{\alpha\beta} = M_{\alpha\beta} = 0$.*

4. Hypersurface $F^{(n-1)}(c)$ of a Finsler space with deformed Randers (α, β) -metric

In this section, we have assumed that the vector field $b_i(x) = \frac{\partial b}{\partial x^i}$ is a gradient of some scalar function $b(x)$ in a deformed Randers (α, β) metric $L(\alpha, \beta) = \alpha^2 + \alpha\beta$. Later, we considered a hypersurface $F^{(n-1)}(c)$ given by the equation $b(x) = c$ [10].

From the parametric equation $x^i = x^i(u^\alpha)$ of $F^{(n-1)}(c)$, we get

$$\frac{\partial b(x)}{\partial u^\alpha} = 0, \quad \frac{\partial b(x)}{\partial x^i} \frac{\partial x^i}{\partial u^\alpha} = 0 \quad \text{and} \quad b_i B_\alpha^i = 0.$$

The above equation shows that $b_i(x)$ is covariant component of a normal vector field of hypersurface $F^{(n-1)}(c)$. Further, we have

$$(4.1) \quad b_i B_\alpha^i = 0, \quad \text{and} \quad b_i y^i = 0 \quad \text{i.e.} \quad \beta = 0,$$

and the induced metric $L(u, v)$ of $F^{(n-1)}(c)$ is given by

$$(4.2) \quad L(u, v) = a_{\alpha\beta} v^\alpha v^\beta, \quad a_{\alpha\beta} = a_{ij} B_\alpha^i B_\beta^j.$$

Writing $\beta = 0$ in the equations (2.3), (2.5) and (2.7) we have

$$(4.3) \quad \begin{cases} p = 2\alpha^2, & q_0 = 0, \\ q_{-1} = \alpha, & q_{-2} = 0, \\ p_0 = \alpha^2, & p_{-1} = 3\alpha, \\ p_{-2} = 4, & \tau = -10\alpha^8(2+b^2)b^2, \\ s_0 = \frac{3}{20\alpha^6(2+b^2)b^2}, \\ s_{-1} = -\frac{3}{10\alpha^7(2+b^2)b^2}, \\ s_{-2} = \frac{5b^2-8}{20\alpha^8b^2(2+b^2)}. \end{cases}$$

From equation (2.6) we have,

$$(4.4) \quad \begin{cases} g^{ij} = \frac{1}{2\alpha^2} a^{ij} - \frac{3}{20\alpha^2 b^2(2+b^2)} b^i b^j + \frac{3}{10\alpha^7 b^2(2+b^2)} (b^i y^j + b^j y^i) \\ \quad - \frac{5b^2-8}{20\alpha^8 b^2(2+b^2)} y^i y^j. \end{cases}$$

Thus along $F^{(n-1)}(c)$, equations (4.3) and (4.4) lead to

$$g^{ij} b_i b_j = \frac{b^2(10b^2+17)}{20\alpha^2(2+b^2)}.$$

So, we get

$$(4.5) \quad b_i(x(u)) = \frac{b}{2\alpha} \sqrt{\frac{10b^2-17}{5(2+b^2)}} N_i,$$

where $b^2 = a^{ij} b_i b_j$, and b is the length of the vector b^i .

Again from the equations (4.4) and (4.5), we get

$$(4.6) \quad b^i = a^{ij} b_j = \frac{20\alpha^2(2+b^2)}{(10b^2+17)} N^i - \frac{6}{\alpha^5(10b^2+17)} y^i.$$

Thus, we have

Theorem 4.1. *The induced Riemannian metric in a Finslerian hypersurface $F^{(n-1)}(c)$ of a Finsler space with deformed Randers (α, β) metric $L = \alpha^2 + \alpha\beta$ is given by (4.2) and the scalar function $b(x)$ is given by (4.5) and (4.6).*

Now the angular metric tensor h_{ij} and metric tensor g_{ij} of F^n are given by

$$(4.7) \quad \begin{cases} h_{ij} = 2\alpha^2 a_{ij} + \alpha(b_i Y_j + b_j Y_i), \\ g_{ij} = 2\alpha^2 a_{ij} + \alpha^2 b_i b_j + 3\alpha(b_i Y_j + b_j Y_i). \end{cases}$$

If $h_{\alpha\beta}^{(a)}$ denotes the angular metric tensor of the Riemannian $a_{ij}(x)$ along Finslerian hypersurfaces $F^{(n-1)}(c)$, then from equations (4.1), (4.7) and (3.2), we have $h_{\alpha\beta} = h_{\alpha\beta}^{(a)}$.

Thus along the Finslerian hypersurface $F^{(n-1)}(c)$, $\frac{\partial p_0}{\partial \beta} = 0$.

Now, using equations (2.3), (2.9), (4.1) and $\frac{\partial p_0}{\partial \beta} = 0$, we get

$$\gamma_1 = 0, \quad m_i = b_i.$$

Again using above result the hv-torsion tensor for the Finslerian hypersurfaces $F^{(n-1)}(c)$ of the Finsler space with deformed Randers (α, β) metric is given by

$$(4.8) \quad C_{ijk} = \frac{3}{4\alpha}(h_{ij}b_k + h_{jk}b_i + h_{ki}b_j).$$

From equations (3.2), (3.3), (3.5), (4.1) and (4.8), we have

$$(4.9) \quad M_{\alpha\beta} = \frac{3b}{8\alpha^2} \sqrt{\frac{10b^2 - 17}{5(2 + b^2)}} h_{\alpha\beta} \quad \text{and} \quad M_\alpha = 0.$$

Therefore, equation (3.6) shows that $H_{\alpha\beta}$ is symmetric. Thus, we have

Theorem 4.2. *The second fundamental v-tensor in a Finslerian hypersurface $F^{(n-1)}(c)$ of a Finsler space with deformed Randers (α, β) metric $L = \alpha^2 + \alpha\beta$ is given by (4.9) and the second fundamental h-tensor $H_{\alpha\beta}$ is symmetric.*

Now from (4.1) we have $b_i B_\alpha^i = 0$. Then we have

$$b_{i|\beta} B_\alpha^i + b_i B_{\alpha|\beta}^i = 0.$$

Therefore, from equation (3.5) and by use of $b_{i|\beta} = b_{i|j} B_\beta^j + b_i|_j N^j H_\beta$, we have

$$(4.10) \quad b_{i|j} B_\alpha^i B_\beta^j + b_{i|j} B_\alpha^i N^j H_\beta + b_i H_{\alpha\beta} N^i = 0.$$

Since $b_i|_j = -b_h C_{ij}^h$, we get

$$b_{i|j} B_\alpha^i N^j = 0.$$

Using above result and equation (4.5) in equation (4.10), we have

$$(4.11) \quad \frac{b}{2\alpha} \sqrt{\frac{10b^2 - 17}{5(2 + b^2)}} H_{\alpha\beta} + b_{i|j} B_{\alpha}^i B_{\beta}^j = 0,$$

because $b_{i|j}$ is symmetric. Now contracting equation (4.11) with v^{β} and using (3.1), we get

$$(4.12) \quad \frac{b}{2\alpha} \sqrt{\frac{10b^2 - 17}{5(2 + b^2)}} H_{\alpha} + b_{i|j} B_{\alpha}^i y^j = 0.$$

Again contracting equation (4.12) by v^{α} and by use of equation (3.1), we have

$$(4.13) \quad \frac{b}{2\alpha} \sqrt{\frac{10b^2 - 17}{5(2 + b^2)}} H_0 + b_{i|j} y^i y^j = 0.$$

From Lemmas 3.1 and 3.2, it is clear that the deformed Randers (α, β) metric of the Finsler hypersurface $F^{(n-1)}(c)$ is a hyperplane of the first kind, if and only if $H_0 = 0$. Thus from equation (4.13) it is obvious that $F^{(n-1)}(c)$ is a hyperplane of the first kind, if and only if $b_{i|j} y^i y^j = 0$. This $b_{i|j}$ being the covariant derivative with respect to CT of F^n defined on y^i , $b_{ij} = \nabla_j b_i$ is the covariant derivative with respect to Riemannian connection $\{^i_{jk}\}$ and is constructed from $a_{ij}(x)$. Hence b_{ij} does not depend on y^i . We shall consider the difference $b_{i|j} - b_{ij}$ where $b_{ij} = \nabla_j b_i$. The difference tensor $D_{jk}^i = \Gamma_{jk}^{*i} - \{^i_{jk}\}$ is given by (2.11). Since b_i is a gradient vector, then from (2.10) we have

$$E_{ij} = b_{ij}, \quad F_{ij} = 0 \quad \text{and} \quad F_j^i = 0.$$

Thus, equation (2.11) reduces to

$$(4.14) \quad \left\{ \begin{array}{l} D_{jk}^i = B^i b_{jk} + B_j^i b_{0k} + B_k^i b_{0j} - b_{0m} g^{im} B_{jk} \\ \quad - C_{jm}^i A_k^m - C_{km}^i A_j^m + C_{jkm} A_s^m g^{is} \\ \quad + \lambda^s (C_{jm}^i C_{sk}^m + C_{km}^i C_{sj}^m - C_{jk}^m C_{ms}^i), \end{array} \right.$$

where

$$(4.15) \quad \left\{ \begin{array}{l} B_i = \alpha^2 b_i + 3\alpha Y_i, \\ B^i = \frac{\alpha^4 b^2 (10b^2 + 19) + 18}{20\alpha^4 b^2 (2 + b^2)} b^i + \frac{3\alpha^6 b^2 (10b^2 + 17) - \alpha^2 (5b^2 - 8) + 6b^2}{20\alpha^7 b^2 (2 + b^2)} y^i, \\ \lambda^m = B^m b_{00}, \\ B_{ij} = \frac{3}{2\alpha} (a_{ij} - \alpha^{-2} Y_i Y_j), \\ B_j^i = -\frac{9}{40\alpha^3 b^2 (b^2 + 2)} b^i b_j, \\ A_k^m = B_k^m b_{00} + B^m b_{k0}. \end{array} \right.$$

In view of equations (4.3) and (4.4), the relation in equation (2.12) becomes possible and by virtue of equation (4.15) we have $B_0^i = 0$, $B_{i0} = 0$ and $A_0^m = B^m b_{00}$.

Now, contracting equation (4.14) by y^k , we get

$$D_{j0}^i = B^i b_{j0} + B_j^i b_{00} - B^m C_{jm}^i b_{00} .$$

Again by contracting the above equation with respect to y^j , we have

$$D_{00}^i = B^i b_{00} = \left\{ \frac{\alpha^4 b^2 (10b^2 + 19) + 18}{20\alpha^4 b^2 (2 + b^2)} b^i + \frac{3\alpha^6 b^2 (10b^2 + 17) - \alpha^2 (5b^2 - 8) + 6b^2}{20\alpha^7 b^2 (2 + b^2)} y^i \right\} b_{00} .$$

Paying attention to equation (4.1), along $F^{(n-1)}(c)$, we get

$$(4.16) \quad \begin{cases} b_i D_{j0}^i = \frac{\alpha^4 b^2 (10b^2 + 19) + 18}{20\alpha^4 (b^2 + 2)} b_{j0} - \frac{9}{40\alpha^3 (b^2 + 2)} b_j b_{00} \\ \quad - \frac{\alpha^4 b^2 (10b^2 + 19) + 18}{20\alpha^4 (b^2 + 2)} b_i b^m C_{jm}^i b_{00} . \end{cases}$$

Contracting (4.16) with y^j implies that

$$(4.17) \quad b_i D_{00}^i = \frac{\alpha^4 b^2 (10b^2 + 19) + 18}{20\alpha^4 (2 + b^2)} .$$

From equations (3.3), (4.5), (4.6), (4.9) and $M_\alpha = 0$, we have

$$b_i b^m C_{jm}^i B_\alpha^j = b^2 M_\alpha = 0 .$$

Thus, the relation of $b_{i|j} = b_{ij} - b_r D_{ij}^r$ the equations (4.16) and (4.17) gives

$$b_{i|j} y^i y^j = b_{00} - b_r D_{00}^r = \frac{\alpha^4 (40 - 10b^4 + 3b^2) - 18}{20\alpha^4 (b^2 + 2)} b_{00} .$$

Consequently equations (4.12) and (4.13) may be written as

$$(4.18) \quad \begin{cases} \frac{b}{2\alpha} \sqrt{\frac{10b^2 - 17}{5(2 + b^2)}} H_\alpha + \frac{\alpha^4 (40 - 10b^4 + 3b^2) - 18}{20\alpha^4 (b^2 + 2)} b_{i|j} B_\alpha^i y^j = 0, \\ \frac{b}{2\alpha} \sqrt{\frac{10b^2 - 17}{5(2 + b^2)}} H_0 + \frac{\alpha^4 (40 - 10b^4 + 3b^2) - 18}{20\alpha^4 (b^2 + 2)} b_{00} = 0. \end{cases}$$

Thus, the condition $H_0 = 0$ is equivalent to $b_{00} = 0$. Using the fact $\beta = b_i y^i = 0$, the condition $b_{00} = 0$ can be written as $b_{ij} y^i y^j = b_i y^i b_j y^j$ for some $c_j(x)$. Thus we can write,

$$(4.19) \quad 2b_{ij} = b_i c_j + b_j c_i .$$

Now from equations (4.1) and (4.19) we get

$$b_{00} = 0, \quad b_{ij} B_\alpha^i B_\beta^j = 0, \quad b_{ij} B_\alpha^i y^j = 0 .$$

Hence from equation (4.18) we get $H_\alpha = 0$, again from equations (4.19) and (4.15) we get $b_{i0} b^i = \frac{c_0 b^2}{2}$, $\lambda^m = 0$, $A_j^i B_\beta^j = 0$ and $B_{ij} B_\alpha^i B_\beta^j = \frac{3}{2\alpha} h_{\alpha\beta}$.

Now with the use of equations (3.3), (4.4), (4.5), (4.6), (4.9) and (4.14) we have

$$(4.20) \quad b_r D_{ij}^r B_\alpha^i B_\beta^j = -\frac{c_0 b^3}{32\alpha^4} \left\{ \frac{(10b^2 - 17)}{5(b^2 + 2)} \right\}^{\frac{3}{2}} h_{\alpha\beta} .$$

Thus the equation (4.11) reduces to

$$(4.21) \quad H_{\alpha\beta} + \frac{3c_0b^2(10b^2 - 17)}{80\alpha^3(b^2 + 2)}h_{\alpha\beta} = 0.$$

Hence the hypersurface $F^{(n-1)}(c)$ is umbilic.

Theorem 4.3. *The necessary and sufficient condition for a Finslerian hypersurface $F^{(n-1)}(c)$ of a Finsler space with Randers deformed (α, β) - metric $L = \alpha^2 + \alpha\beta$ to be a hyperplane of the first kind is (4.19).*

Corollary 4.4. *The second fundamental h-tensor in a Finslerian hypersurface $F^{(n-1)}(c)$ of a Finsler space with Randers deformed (α, β) - metric $L = \alpha^2 + \alpha\beta$ is directly proportional to its angular metric tensor.*

Now from Lemma 3.3, $F^{(n-1)}(c)$ is a hyperplane of the second kind if and only if $H_\alpha = 0$ and $H_{\alpha\beta} = 0$. Thus from (4.20), we get

$$c_0 = c_i(x)y^i = 0.$$

Now, there exists a function $\psi(x)$ such that

$$c_i(x) = \psi(x)b_i(x).$$

Therefore, from equation (4.19), we get

$$2b_{ij} = b_i(x)\psi(x)b_j(x) + b_j(x)\psi(x)b_i(x).$$

This can also be written as

$$b_{ij} = \psi(x)b_ib_j.$$

Theorem 4.5. *The necessary and sufficient condition for a Finslerian hypersurface $F^{(n-1)}(c)$ of a Finsler space with Randers deformed (α, β) - metric $L = \alpha^2 + \alpha\beta$ to be a hyperplane of the second kind is (4.21).*

Again Lemma 3.4, together with (4.9) and $M_\alpha = 0$ shows that $F^{(n-1)}(c)$ is not a hyperplane of the third kind.

Theorem 4.6. *The Finslerian hypersurface $F^{(n-1)}(c)$ of a Finsler space with Randers deformed (α, β) - metric $L = \alpha^2 + \alpha\beta$ is not a hyperplane of the third kind.*

5. Conclusion

In the present paper, we have obtained the condition under which a deformed Randers (α, β) -metric which is defined by equation (1.1) of a 2-degree homogeneous function and this metric also formed a nonholonomic frame of a Finsler space [12]. Further, we have obtained the Theorems 4.3 and 4.5 and they state that the condition under which the Finslerian hypersurfaces of this metric is a hyperplane of the first and the second kind. Again, we have proved in Theorem 4.6 that it is not a hyperplane of the third kind.

Since the noholonomic frame is defined for the Finsler metric of two-degree homogeneous function and this work can also be extended for the study of Finslerian hypersurfaces for those Finsler metric which framed nonholonomic frame for future studies.

References

- [1] CHAUBEY, V. K., AND MISHRA, A. Hypersurfaces of a Finsler space with a special (α, β) -metric. *Izv. Nats. Akad. Nauk Armenii Mat.* 52, 1 (2017), 38–46.
- [2] CHAUBEY, V. K., AND TRIPATHI, B. K. Finslerian hypersurfaces of a Finsler spaces with (γ, β) -metric. *J. Dyn. Syst. Geom. Theor.* 12, 1 (2014), 19–27.
- [3] KITAYAMA, M. On Finslerian hypersurfaces given by β -changes. *Balkan J. Geom. Appl.* 7, 2 (2002), 49–55.
- [4] KUMAR, P., NARASIMHAMURTHY, S. K., NAGARAJA, H. G., AND AVEESH, S. T. On a special hypersurface of a Finsler space with (α, β) -metric. *Tbil. Math. J.* 2 (2009), 51–60.
- [5] LEE, I. Y., PARK, H. Y., AND LEE, Y. D. On a hypersurface of a special finsler spaces with a metric $(\alpha + \frac{\beta^2}{\alpha})$. *Korean J. Math. Sciences* (2001), 93 – 101.
- [6] MATSUMOTO, M. The induced and intrinsic Finsler connections of a hypersurface and Finslerien projective geometry. *J. Math. Kyoto Univ.* 25, 1 (1985), 107–144.
- [7] MATSUMOTO, M. Theory of Finsler spaces with (α, β) -metric. *Rep. Math. Phys.* 31, 1 (1992), 43–83.
- [8] PANDEY, T. N., AND TRIPATHI, B. K. On a hypersurface of a Finsler space with special (α, β) -metric. *Tensor (N.S.)* 68, 2 (2007), 158–166.
- [9] SHIBATA, C. On Finsler spaces with an (α, β) -metric. *J. Hokkaido Univ. Ed. Sect. II A* 35, 1 (1984), 1–16.
- [10] SINGH, U. P., AND KUMARI, B. On a hypersurface of a Matsumoto space. *Indian J. Pure Appl. Math.* 32, 4 (2001), 521–531.
- [11] TRIPATHI, B. K. Hypersurfaces of a finsler space with deformed berwal-infinite series metric. *TWMS Journal of Applied and Engineering Mathematics*, 2 (2020), 296–304.
- [12] TRIPATHI, B. K., AND CHAUBEY, V. Nonholonomic Frames for Finsler Spaces with special (α, β) -metric. *J. Math. Comput. Sci.* 8, 2 (2018), 290–296.

Received by the editors June 3, 2021

First published online January 11, 2022