

# Mathias–Příkrý forcing and generic ultrafilters

David Chodounský

Institute of Mathematics CAS

# Ultrafilters on $\omega$

## Definition

An ultrafilter  $\mathcal{F}$  on  $\omega$  is a *P-point* if for each  $\mathcal{C} \in [\mathcal{F}]^\omega$  there is a pseudo-intersection  $P \in \mathcal{F}$  such that  $P \subset^* F$  for each  $F \in \mathcal{C}$ .

## Definition

An ultrafilter  $\mathcal{F}$  on  $\omega$  is *selective* if for each  $\{A_i : i \in \omega\}$ , a partition of  $\omega$  disjoint with  $\mathcal{F}$  there is a selector  $S \in \mathcal{F}$  such that  $|S \cap A_i| = 1$  for each  $i \in \omega$ .

# Ultrafilters on $\omega$

## Definition

An ultrafilter  $\mathcal{F}$  on  $\omega$  is a *P-point* if for each  $\mathcal{C} \in [\mathcal{F}]^\omega$  there is a pseudo-intersection  $P \in \mathcal{F}$  such that  $P \subset^* F$  for each  $F \in \mathcal{C}$ .

## Definition

An ultrafilter  $\mathcal{F}$  on  $\omega$  is *selective* if for each  $\{A_i : i \in \omega\}$ , a partition of  $\omega$  disjoint with  $\mathcal{F}$  there is a selector  $S \in \mathcal{F}$  such that  $|S \cap A_i| = 1$  for each  $i \in \omega$ .

Let  $\mathcal{F}$  be an ultrafilter in  $\omega$ . The following properties are equivalent:

1.  $\mathcal{F}$  is selective,
2. For each  $c: [\omega]^2 \rightarrow 2$  there exists a  $c$ -homogeneous set  $F \in \mathcal{F}$ ,

# Ultrafilters on $\omega$

## Definition

An ultrafilter  $\mathcal{F}$  on  $\omega$  is a *P-point* if for each  $\mathcal{C} \in [\mathcal{F}]^\omega$  there is a pseudo-intersection  $P \in \mathcal{F}$  such that  $P \subset^* F$  for each  $F \in \mathcal{C}$ .

## Definition

An ultrafilter  $\mathcal{F}$  on  $\omega$  is *selective* if for each  $\{A_i : i \in \omega\}$ , a partition of  $\omega$  disjoint with  $\mathcal{F}$  there is a selector  $S \in \mathcal{F}$  such that  $|S \cap A_i| = 1$  for each  $i \in \omega$ .

Let  $\mathcal{F}$  be an ultrafilter in  $\omega$ . The following properties are equivalent:

1.  $\mathcal{F}$  is selective,
2. For each  $c: [\omega]^2 \rightarrow 2$  there exists a  $c$ -homogeneous set  $F \in \mathcal{F}$ ,
3.  $\mathcal{F} \cap \mathcal{I} \neq \emptyset$  for each tall analytic ideal  $\mathcal{I}$ .

# Ultrafilters on $\omega$

## Definition

An ultrafilter  $\mathcal{F}$  on  $\omega$  is a *P-point* if for each  $\mathcal{C} \in [\mathcal{F}]^\omega$  there is a pseudo-intersection  $P \in \mathcal{F}$  such that  $P \subset^* F$  for each  $F \in \mathcal{C}$ .

## Definition

An ultrafilter  $\mathcal{F}$  on  $\omega$  is *selective* if for each  $\{A_i: i \in \omega\}$ , a partition of  $\omega$  disjoint with  $\mathcal{F}$  there is a selector  $S \in \mathcal{F}$  such that  $|S \cap A_i| = 1$  for each  $i \in \omega$ .

Let  $\mathcal{F}$  be an ultrafilter in  $\omega$ . The following properties are equivalent:

1.  $\mathcal{F}$  is selective,
2. For each  $c: [\omega]^2 \rightarrow 2$  there exists a  $c$ -homogeneous set  $F \in \mathcal{F}$ ,
3.  $\mathcal{F} \cap \mathcal{I} \neq \emptyset$  for each tall analytic ideal  $\mathcal{I}$ .

## Theorem (Zapletal)

An ultrafilter  $\mathcal{F}$  is a *P-point* iff for each analytic ideal  $\mathcal{I} \subset \mathcal{F}^*$  there is an  $F_\sigma$  ideal  $\mathcal{C}$  such that  $\mathcal{I} \subseteq \mathcal{C} \subseteq \mathcal{F}^*$ .

# Generic filters

## Theorem

*The generic filter on the poset  $(\mathcal{P}(\omega) \setminus \text{Fin}, \subset^*)$  is a selective ultrafilter.*

# Generic filters

## Theorem

*The generic filter on the poset  $(\mathcal{P}(\omega) \setminus \text{Fin}, \subset^*)$  is a selective ultrafilter.*

## Theorem (Todorćević)

*(LC)<sup>1</sup> An ultrafilter is selective if and only if it is a generic filter on  $\mathcal{P}(\omega) \setminus \text{Fin}$  over  $L(\mathbb{R})$ .*

---

<sup>1</sup>(LC) denotes the assumption that there exist sufficiently large cardinals in  $V$ .

# Generic filters

## Theorem

*The generic filter on the poset  $(\mathcal{P}(\omega) \setminus \text{Fin}, \subset^*)$  is a selective ultrafilter.*

## Theorem (Todorćević)

*(LC)<sup>1</sup> An ultrafilter is selective if and only if it is a generic filter on  $\mathcal{P}(\omega) \setminus \text{Fin}$  over  $L(\mathbb{R})$ .*

Let  $\mathcal{I}$  be an  $F_\sigma$  ideal on  $\omega$ . Denote by  $\mathbb{Q}_{\mathcal{I}}$  the forcing  $(\mathcal{P}(\omega) \setminus \mathcal{I}, \subset^*)$ .

---

<sup>1</sup>(LC) denotes the assumption that there exist sufficiently large cardinals in  $V$ .



# Generic filters

## Theorem

*The generic filter on the poset  $(\mathcal{P}(\omega) \setminus \text{Fin}, \subset^*)$  is a selective ultrafilter.*

## Theorem (Todorcevic)

*(LC)<sup>1</sup> An ultrafilter is selective if and only if it is a generic filter on  $\mathcal{P}(\omega) \setminus \text{Fin}$  over  $L(\mathbb{R})$ .*

Let  $\mathcal{I}$  be an  $F_\sigma$  ideal on  $\omega$ . Denote by  $\mathbb{Q}_{\mathcal{I}}$  the forcing  $(\mathcal{P}(\omega) \setminus \mathcal{I}, \subset^*)$ .

## Theorem (Zapletal, Ch.)

*(LC)  $\mathcal{F}$  is a  $\mathbb{Q}_{\mathcal{I}}$ -generic filter over  $L(\mathbb{R})$  iff*

- 1.  $\mathcal{F}$  is a  $P$ -point disjoint with  $\mathcal{I}$ , and*
- 2. for each closed set  $C \subset \mathcal{P}(\omega)$  disjoint with  $\mathcal{F}$  there is  $e \in \mathcal{F}^*$  such that  $C \subseteq \langle \mathcal{I}, \{e\} \rangle$ .*

---

<sup>1</sup>(LC) denotes the assumption that there exist sufficiently large cardinals in  $V$ .

## Definition (Mathias forcing)

$$\mathbb{M} = \{\langle a, F \rangle : a \in [\omega]^{<\omega}, F \in [\omega]^\omega\}$$

$$\langle a, F \rangle < \langle b, H \rangle \quad \text{if} \quad b \sqsubseteq a, F \subset H, \text{ and } a \setminus b \subset H.$$

## Definition (Mathias–Příkrý forcing)

Let  $\mathcal{F}$  be a filter on  $\omega$ .

$$\mathbb{M}(\mathcal{F}) = \{\langle a, F \rangle : a \in [\omega]^{<\omega}, F \in \mathcal{F}\}$$

$$\langle a, F \rangle < \langle b, H \rangle \quad \text{if} \quad b \sqsubseteq a, F \subset H, \text{ and } a \setminus b \subset H.$$

## Definition (Mathias–Příkrý forcing)

Let  $\mathcal{F}$  be a filter on  $\omega$ .

$$\mathbb{M}(\mathcal{F}) = \{\langle a, F \rangle : a \in [\omega]^{<\omega}, F \in \mathcal{F}\}$$

$$\langle a, F \rangle < \langle b, H \rangle \quad \text{if} \quad b \sqsubseteq a, F \subset H, \text{ and } a \setminus b \subset H.$$

## Definition (Mathias real for $\mathcal{F}$ )

$x = \bigcup \{a : \langle a, F \rangle\} \in \mathbb{G}$ , where  $\mathbb{G}$  is an  $\mathbb{M}(\mathcal{F})$  generic filter.

## Fact

*A Mathias real is a pseudo-intersection of  $\mathcal{F}$  ( $x \subseteq^* F$  for each  $F \in \mathcal{F}$ ).*

## Definition (Mathias–Příkrý forcing)

Let  $\mathcal{F}$  be a filter on  $\omega$ .

$$\mathbb{M}(\mathcal{F}) = \{\langle a, F \rangle : a \in [\omega]^{<\omega}, F \in \mathcal{F}\}$$

$$\langle a, F \rangle < \langle b, H \rangle \quad \text{if} \quad b \sqsubseteq a, F \subset H, \text{ and } a \setminus b \subset H.$$

## Definition (Mathias real for $\mathcal{F}$ )

$x = \bigcup \{a : \langle a, F \rangle\} \in \mathbb{G}$ , where  $\mathbb{G}$  is an  $\mathbb{M}(\mathcal{F})$  generic filter.

## Fact

*A Mathias real is a pseudo-intersection of  $\mathcal{F}$  ( $x \subseteq^* F$  for each  $F \in \mathcal{F}$ ).*

## Definition

$U \subset [\omega]^{<\omega}$  is an  $\mathcal{F}$ -universal set if  $[F]^{<\omega} \cap U \neq \emptyset$  for each  $F \in \mathcal{F}$ .

## Fact

$[x]^{<\omega} \cap U \neq \emptyset$  for each  $\mathcal{F}$ -universal set  $U$ .

## Definition (Pseudogeneric real for $\mathcal{F}$ )

Let  $\mathcal{F}$  be a filter on  $\omega$ . A set  $m \subset \omega$  is a *pseudogeneric real for  $\mathcal{F}$*  if

1.  $m \subseteq^* F$  for each  $F \in \mathcal{F}$ ,
2.  $[m]^{<\omega} \cap U \neq \emptyset$  for each  $\mathcal{F}$ -universal set  $U$ .

## Definition (Pseudogeneric real for $\mathcal{F}$ )

Let  $\mathcal{F}$  be a filter on  $\omega$ . A set  $m \subset \omega$  is a *pseudogeneric real* for  $\mathcal{F}$  if

1.  $m \subseteq^* F$  for each  $F \in \mathcal{F}$ ,
2.  $[m]^{<\omega} \cap U \neq \emptyset$  for each  $\mathcal{F}$ -universal set  $U$ .

## Theorem

If  $m \subset \omega$  is a pseudogeneric real for  $\mathcal{F}$  and  $c \subset \omega$  is a Cohen real, then  $m \cap c$  is a Mathias real for  $\mathcal{F}$ .

## Theorem

(LC)  $\mathcal{F}$  is a  $\mathbb{Q}_{\mathcal{I}}$ -generic filter over  $L(\mathbb{R})$  iff

1.  $\mathcal{F}$  is a  $P$ -point disjoint with  $\mathcal{I}$ , and
2. for each closed set  $C \subset \mathcal{P}(\omega)$  disjoint with  $\mathcal{F}$  there is  $e \in \mathcal{F}^*$  such that  $C \subseteq \langle \mathcal{I}, \{e\} \rangle$ .



## Theorem

(LC)  $\mathcal{F}$  is a  $\mathbb{Q}_{\mathcal{I}}$ -generic filter over  $L(\mathbb{R})$  iff

1.  $\mathcal{F}$  is a  $P$ -point disjoint with  $\mathcal{I}$ , and
2. for each closed set  $C \subset \mathcal{P}(\omega)$  disjoint with  $\mathcal{F}$  there is  $e \in \mathcal{F}^*$  such that  $C \subseteq \langle \mathcal{I}, \{e\} \rangle$ .

## Lemma

Let  $\mathcal{U}$  be a  $P$ -point. Assume there is (in some extension of  $V$ ) an elementary embedding  $j: V \rightarrow M$  such that  $\mathbb{R} \cap V$  is countable in  $M$ . Then there is a Mathias real  $g \in j(\mathcal{F})$  (over  $V$ ).

## Theorem

(LC)  $\mathcal{F}$  is a  $\mathbb{Q}_{\mathcal{I}}$ -generic filter over  $L(\mathbb{R})$  iff

1.  $\mathcal{F}$  is a  $P$ -point disjoint with  $\mathcal{I}$ , and
2. for each closed set  $C \subset \mathcal{P}(\omega)$  disjoint with  $\mathcal{F}$  there is  $e \in \mathcal{F}^*$  such that  $C \subseteq \langle \mathcal{I}, \{e\} \rangle$ .

## Lemma

Let  $\mathcal{U}$  be a  $P$ -point. Assume there is (in some extension of  $V$ ) an elementary embedding  $j: V \rightarrow M$  such that  $\mathbb{R} \cap V$  is countable in  $M$ . Then there is a Mathias real  $g \in j(\mathcal{F})$  (over  $V$ ).

## Lemma

Let  $\mathcal{F}$  be as in the theorem. Suppose  $D \in L(\mathbb{R})$  is open dense in  $\mathbb{Q}_{\mathcal{I}}$ . Then  $M(\mathcal{F}) \Vdash \dot{g} \in D^{V[\dot{g}]}$ .

## Lemma

Let  $\mathcal{U}$  be a  $P$ -point. Assume there is (in some extension of  $V$ ) an elementary embedding  $j: V \rightarrow M$  such that  $\mathbb{R} \cap V$  is countable in  $M$ . Then there is a Mathias real  $g \in j(\mathcal{F})$  (over  $V$ ).

## Lemma

Let  $\mathcal{F}$  be as in the theorem. Suppose  $D \in L(\mathbb{R})$  is open dense in  $\mathbb{Q}_{\mathcal{I}}$ . Then  $M(\mathcal{F}) \Vdash \dot{g} \in D^{V[\dot{g}]}$ .

## Lemma

Let  $\mathcal{U}$  be a  $P$ -point. Assume there is (in some extension of  $V$ ) an elementary embedding  $j: V \rightarrow M$  such that  $\mathbb{R} \cap V$  is countable in  $M$ . Then there is a Mathias real  $g \in j(\mathcal{F})$  (over  $V$ ).

## Lemma

Let  $\mathcal{F}$  be as in the theorem. Suppose  $D \in L(\mathbb{R})$  is open dense in  $\mathbb{Q}_{\mathcal{I}}$ . Then  $M(\mathcal{F}) \Vdash \dot{g} \in D^{V[\dot{g}]}$ .

Suppose  $D \in L(\mathbb{R})$  is open dense in  $\mathbb{Q}_{\mathcal{I}}$ . We need to show that  $D \cap \mathcal{F} \neq \emptyset$ .

## Lemma

Let  $\mathcal{U}$  be a  $P$ -point. Assume there is (in some extension of  $V$ ) an elementary embedding  $j: V \rightarrow M$  such that  $\mathbb{R} \cap V$  is countable in  $M$ . Then there is a Mathias real  $g \in j(\mathcal{F})$  (over  $V$ ).

## Lemma

Let  $\mathcal{F}$  be as in the theorem. Suppose  $D \in L(\mathbb{R})$  is open dense in  $\mathbb{Q}_{\mathcal{I}}$ . Then  $M(\mathcal{F}) \Vdash \dot{g} \in D^{V[\dot{g}]}$ .

Suppose  $D \in L(\mathbb{R})$  is open dense in  $\mathbb{Q}_{\mathcal{I}}$ . We need to show that  $D \cap \mathcal{F} \neq \emptyset$ .

Pass to an extension  $V[G]$  where  $j: V \rightarrow M$  exists.

## Lemma

Let  $\mathcal{U}$  be a  $P$ -point. Assume there is (in some extension of  $V$ ) an elementary embedding  $j: V \rightarrow M$  such that  $\mathbb{R} \cap V$  is countable in  $M$ . Then there is a Mathias real  $g \in j(\mathcal{F})$  (over  $V$ ).

## Lemma

Let  $\mathcal{F}$  be as in the theorem. Suppose  $D \in L(\mathbb{R})$  is open dense in  $\mathbb{Q}_I$ . Then  $M(\mathcal{F}) \Vdash \dot{g} \in D^{V[\dot{g}]}$ .

Suppose  $D \in L(\mathbb{R})$  is open dense in  $\mathbb{Q}_I$ . We need to show that  $D \cap \mathcal{F} \neq \emptyset$ .

Pass to an extension  $V[G]$  where  $j: V \rightarrow M$  exists. There is a Mathias real  $g \in j(\mathcal{F})$ .

## Lemma

Let  $\mathcal{U}$  be a  $P$ -point. Assume there is (in some extension of  $V$ ) an elementary embedding  $j: V \rightarrow M$  such that  $\mathbb{R} \cap V$  is countable in  $M$ . Then there is a Mathias real  $g \in j(\mathcal{F})$  (over  $V$ ).

## Lemma

Let  $\mathcal{F}$  be as in the theorem. Suppose  $D \in L(\mathbb{R})$  is open dense in  $\mathbb{Q}_I$ . Then  $M(\mathcal{F}) \Vdash \dot{g} \in D^{V[\dot{g}]}$ .

Suppose  $D \in L(\mathbb{R})$  is open dense in  $\mathbb{Q}_I$ . We need to show that  $D \cap \mathcal{F} \neq \emptyset$ .

Pass to an extension  $V[G]$  where  $j: V \rightarrow M$  exists.

There is a Mathias real  $g \in j(\mathcal{F})$ .

Now  $g \in D^{V[G]}$ , i.e.  $g \in j(D) \cap j(\mathcal{F})$ .